

**July 2009
Edition 1**

NZAF HIV Prevention Plan 2009-2014

Mahere Ārai Mate Āraikore



NEW ZEALAND AIDS FOUNDATION
Te Tūāpapa Mate Āraikore o Aotearoa

Contents

Rārangi Upoko

Introduction : He Kupu Whakataki	5
Process : Tikanga	6
HIV In Aotearoa New Zealand : Mate Āraikore Aotearoa	7
Background And Context : Horopaki	11
The Plan : Te Mahere	15
Goal One : Whāinga Tuatahi: Increase rates of condom use for anal sex between men	17
Goal Two : Whāinga Tuarua: Increase STI and HIV testing rates for gay and bisexual men	21
Goal Three : Whāinga Tuatoru: Increase rates of condom use for first anal sex between men	22
Goal Four : Whāinga Tuawha: Increase rates of condom use within New Zealand-based African communities	23
Programme Logic Model : Aromātai Ture	
Programme Evaluation Framework : Aromātai Hātaka	24
Reference List : Rārangi Pukapuka	29

Introduction

He Kupu Whakataki

The New Zealand AIDS Foundation (NZAF) aims to achieve its vision of a world without HIV and AIDS through its mission of preventing the transmission of HIV and supporting those affected by HIV to maximise their health and wellbeing. In the 24 years of the HIV epidemic in Aotearoa New Zealand, the landscape the NZAF works in has changed. Homosexual Law Reform, the internet and the advent of antiretroviral therapies are a few of the significant events that have changed the environment that the NZAF first confronted in 1984.

This plan marks the first comprehensive review of NZAF HIV prevention programme delivery for many years. In 2008, New Zealand recorded the highest number of new HIV diagnoses ever, and with the continued rise in new diagnoses since 2002, it is both timely and necessary for this review to occur.

The NZAF HIV Prevention Plan 2009-14 is knowledge based: robust scientific evidence and the best epidemiological and behavioural surveillance available provided the basis for this plan. This evidence has unequivocally shown that to prevent HIV transmission, prevention initiatives must occur in the places where HIV prevalence is, and those initiatives must be focused on changing the risk behaviours of the groups most at risk. To respond effectively to the changing environment of the epidemic the NZAF must also embrace social marketing and new technology while maintaining the values and principles of the communities we serve, including a clear and sincere commitment to takatāpui and whānau.

Increasing condom use is the core tenet of the plan because using condoms and lube remains the single most effective intervention to prevent the transmission of HIV.

This plan recognises that the people most at risk of HIV are influenced by their partners, whānau, family, friends, colleagues, employers, communities, and the society in which they live. To reduce HIV prevalence, Aotearoa New Zealand must foster a condom culture; a social movement that promotes, supports and demands consistent use of condoms and lube.

The NZAF HIV Prevention Plan 2009-14 and the new directions it heralds are singular in purpose – they will ensure that the NZAF reduces the rate of HIV transmission in Aotearoa New Zealand.



Simon Harger-Forde
Director HIV Prevention and Communications



Rachael Le Mesurier
Executive Director

July 2009

Process Tikanga

A process that involved close analysis of scientific evidence and surveillance literature has been followed to develop this plan.

The NZAF identified a five stage process to develop this plan. In stage one a group of NZAF staff analysed literature identified as appropriate and useful to develop HIV prevention and intervention programmes. The research included behavioural surveillance data on gay men in New Zealand (Gay Auckland Periodic Sex Survey and Gay Online Sex Survey), HIV epidemiology from the AIDS Epidemiology Group at the University of Otago, data from Youth 2007 (Adolescent Health Research Group, University of Auckland), a specific study into non condom use in Auckland (Massey University), data from HIV Futures NZ² (Latrobe University), and a variety of international HIV prevention literature.

The question 'what do we know?' was asked in an externally facilitated discussion and then the information collated was extrapolated to identify key target audiences.

These audiences were then considered by a smaller group of people (stage two), and four key behaviours were identified as the most effective to reduce transmission of HIV in Aotearoa New Zealand. Several health outcomes were proposed for each behaviour and collated into a simple programme logic model. The initial group then reconvened to integrate feedback from the external reviewers of stage two and to develop the programme logic model (stage three). The programme logic model identified specific activities, ensuring that the outcomes proposed matched the goals of the NZAF Strategic Plan 2005-2010 and the service specifications in the existing Ministry of Health contract, and that the activities effectively targeted the groups identified at the beginning of the process.

Stage four focused on detailed organisational planning, including budgeting for the new model and identifying a structure and specific staff roles. Stage five focused on the implementation of the plan and involved organisational change, engaging with key external stakeholders and establishing a reference group of key stakeholders to guide the implementation.


NZAF staff were engaged in this process and had the opportunity to participate in the process.

HIV in Aotearoa New Zealand Mate Āraikore Aotearoa

Since 1985, AIDS has been notifiable (by anonymous coding) in New Zealand. HIV is not notifiable (as at June 2009) but a voluntary system of anonymous surveillance has been in place since 1986. Since the mid-1990s, AIDS deaths have decreased due to the availability of antiretroviral therapies. Antiretroviral therapies delay the progression of HIV infection to AIDS. The decrease in AIDS diagnoses has led to AIDS being less useful as a means of tracking the HIV epidemic. Since 1996, an enhanced surveillance system for HIV diagnoses has provided detailed information on HIV diagnoses and improved understanding of patterns of HIV infection (Saxton, Dickson & Hughes, 2006).

AIDS diagnoses peaked in 1989 (see Figure 1). New Zealand was one of the first countries in the world to experience a decline in AIDS incidence (Sharples, et al, 1996), and the major factors responsible for this are likely to have been the reduction in HIV infection among men who have sex with men (MSM) in the late 1980s, and the effective prevention of epidemics in other at risk populations (sex workers and injecting drug users (IDU)).


Figure 1



Annual number of diagnoses of AIDS and deaths among people notified with AIDS
(The number of notifications and deaths for 2007 might rise due to delayed reports).
(Source: AIDS Epidemiology Group, University of Otago, 2008)

HIV tests may not be carried out until years after HIV has been acquired so the number of HIV diagnoses should not be interpreted as the number of people annually infected with HIV. In addition, not all people at risk will have been tested.

Exposure categories of annual HIV diagnoses via Western blot antibody testing.
(Note: Does not distinguish between infections acquired in NZ and overseas).



Data provided by AIDS Epidemiology Group, Department of Preventive and Social Medicine, University of Otago.
Graph produced by Research, Analysis and Information Unit, New Zealand AIDS Foundation; 2008.

The HIV epidemic in New Zealand is comprised of two distinct sub-epidemics, one among MSM and one among heterosexual migrants.


Throughout the past decade (1999-2008) the majority of MSM were infected in New Zealand. Since 2002 the number of infections acquired in New Zealand has risen while the number acquired overseas has remained level. In most years over the past decade, there have been more diagnoses of HIV among MSM in Auckland than in the rest of New Zealand combined.

The number of people diagnosed with HIV acquired through heterosexual contact peaked in 2006, probably a reflection of legislative changes requiring more HIV testing for immigration purposes in November 2005. In contrast to the situation with MSM, the majority of people with heterosexually acquired HIV over the period 1999-2008 were infected overseas; 81% in 2008. These infections were mainly acquired in countries where there is a high prevalence of HIV in the heterosexual population.


Perinatal infection (HIV transmission from mother to child) primarily occurs in another country and is later diagnosed in New Zealand. Perinatal HIV transmission has not occurred in NZ, by a woman who knew she was HIV positive, since 1995.

In 2008, 184 people were newly diagnosed with HIV in New Zealand through antibody testing. This is one more than the previous high of 183 new diagnoses in 2005. 91 were men infected through sex with other men, 61 (39 men and 22 women) through heterosexual contact, 2 through injecting drug use, 2 through a transfusion (overseas), 4 were children infected through mother-to-child transmission (3 overseas and 1 in New Zealand), 3 people had another means of infection, and for 21 people the means of infection was unknown or unreported.

Over the five years from 2004 to 2008 inclusive, MSM have accounted for 79% of the total HIV diagnoses by antibody testing in New Zealand, where HIV infection has occurred in New Zealand.



Place of infection of MSM diagnosed by antibody test, by year of diagnosis
(Source: AIDS Epidemiology Group, AIDS-New Zealand, 2009)



Place of infection of those infected through heterosexual contact, diagnosed by antibody test, by year of diagnosis
(Source: AIDS Epidemiology Group, AIDS-New Zealand, 2009)

Ethnicity* by time of diagnosis in New Zealand for those found to be infected with HIV by antibody test and first viral load test.
(A small number of transsexuals are included with the males).
(Source: AIDS Epidemiology Group, AIDS-New Zealand, 2009)

Ethnicity	Sex	HIV Infection*									
		1996-1999		2000-2004		2005-2007		2008		Total	
		No.	%	No.	%	No.	%	No.	%	No.	%
European/Pakeha	Male	257	53.5	337	46.4	250	40.5	97	42.7	941	45.8
	Female	25	5.2	30	4.1	21	3.4	7	3.1	83	4.0
Maori†	Male	29	6.0	41	5.6	46	7.4	17	7.5	133	6.5
	Female	4	0.8	5	0.7	5	0.8	1	0.4	15	0.7
Pacific Island	Male	4	0.8	19	2.6	16	2.6	4	1.8	43	2.1
	Female	4	0.8	10	1.4	7	1.1	2	0.9	23	1.1
African	Male	55	11.4	70	9.6	77	12.5	21	9.2	223	10.9
	Female	33	6.9	85	11.6	81	13.1	18	7.9	217	10.6
Asian	Male	32	6.7	69	9.5	52	8.4	19	8.3	172	8.4
	Female	16	3.3	32	4.4	19	3.1	5	2.2	72	3.5
Other	Male	7	1.4	16	2.2	17	2.7	13	5.7	53	2.6
	Female	0	0.0	2	0.3	7	1.1	1	0.4	10	0.5
Awaiting information/undetermined	Male	13	2.7	13	1.8	12	1.9	19	8.3	57	2.8
	Female	1	0.2	0	0.0	7	1.1	3	1.3	11	0.5
TOTAL		480	100.0	729	100.0	617	100.0	227	100.0	2053	100.0

* Includes people who have developed AIDS. HIV numbers are recorded by time of diagnosis for those reported through antibody testing and by time of first viral load for those reported through viral load testing. The latter include many who have initially been diagnosed overseas and not had an antibody test here. The date of initial diagnosis may have preceded the viral load date by months or years.
† Information on ethnicity of people diagnosed with HIV only collected since 1996.
‡ Includes people who belong to Maori and another ethnic group.

Background and Context Horopaki

The NZAF and the Ottawa Charter

The NZAF has a commitment to the Ottawa Charter for Health Promotion, and like many public health organisations across the world, the NZAF has developed its HIV prevention work using the five action areas of health promotion identified within the charter. The NZAF can align much of its prevention work since 1985 to this model of health promotion:

1. Building healthy public policy – the NZAF recognised early in its history that advocating for changes to legislation and public policy that discriminated against gay and bisexual men, sex workers, injecting drug users, and people living with HIV was a critical step in the fight against HIV and AIDS.
2. Strengthening community action – the NZAF has contributed to the development of gay communities and African communities across New Zealand in the form of practical assistance (venues, funding) and facilitation, recognising strong communities as protective factors against HIV.
3. Creating supportive environments – the NZAF has put significant energy into organising, promoting and advocating for venues, events, social and leisure groups that contribute to healthy social environments for gay and bisexual men and African communities.
4. Developing personal skills – over many years the NZAF has provided workshops for gay and bisexual men, young people and people from African communities and has developed and distributed significant amounts of sexual health and HIV information through a range of mediums.
5. Re-orientating health care services toward prevention of illness and promotion of health – the NZAF has worked to support and train professionals working with the communities most at risk of HIV, has worked to provide new and innovative services for gay men and advocated for the reorientation of services provided by others to gay and bisexual men.

The NZAF has not reviewed this overall approach to the delivery of HIV prevention programmes since 2001. As new funding or ideas emerged they were added to the current programme without rethinking the programme's content and approach.

Commitment to Tangata Whenua: Takatāpui and Whānau

The NZAF is committed to honouring the spirit of Te Tiriti o Waitangi through a practical commitment to biculturalism. For the past thirteen years this has included a parallel model of bicultural development for HIV prevention focused on takatāpui tane.

Unlike many health issues in Aotearoa New Zealand, for example sexually transmitted infections (STIs), there is no significant inequality between Māori and non-Māori in terms of HIV transmission. 155 Māori (137 men and 18 women) have been diagnosed with HIV since 1998, compared to 1,069 Pākehā (982 men and 87 women). The total number of HIV diagnoses since 1998 is 2,146, therefore 7.2% of diagnoses were Māori.

Māori are also not under represented in HIV testing or over represented in AIDS diagnosis statistics, which would suggest that it is unlikely there is a significantly higher level of undiagnosed HIV within Māori populations. The NZAF HIV Prevention Plan 2009-14 ensures a continued commitment to appropriate and effective interventions for Māori by targeting takatāpui and whānau. Organisational strengthening measures will ensure staff with strong skills in Māori public health, especially in relation to takatāpui health are in key programme management, social marketing and community engagement roles.

Changing Epidemic – Changing Communities

As the HIV epidemic has changed, so have the communities it affects. New generations of gay and bisexual men live in vastly different social and political environments than the men who first responded to HIV.

Earlier generations of gay and bisexual men were part of a subculture united in part, by shared criminality. Homosexual behaviour was seen by many as rebellion against staid norms and unjust laws. Some men created an underground, outlaw culture based on desire.

In the decades after 1970 many gay men were increasing politicised as part of a wider sexual liberation movement. Over time the goal became less radical and now in developed countries the movement has generally become a 'rights' movement.

Today, homosexual and bisexual behaviour between men is no longer illegal in New Zealand; gay and bisexual men have human rights and can enter into legally sanctioned civil unions. This means that young men will experience their sexuality – and the HIV epidemic – very differently to men of earlier generations.

Socially the young gay couple has little reason to see themselves as being very different from any other young couple and want what is perceived as the same freedoms.

Today, it is a far more complicated world than that in which HIV first emerged. Segmentation of the audiences is vital. The audiences now include the 'mainstream' and the 'significant others' of gay and bisexual men who influence and support behaviour change.

Changing Epidemic – Changing Direction

Some commentators suggest that we are in a time that is 'post health promotion'. While this discourse poses interesting opportunity for debate, the NZAF HIV Prevention Plan 2009-14 does not support the 'disposal of health promotion', but does seek to respond to the HIV epidemic as it is now in New Zealand, and to use new and innovative approaches. Many of the activities suggested in the new programme are easily aligned to the Ottawa Charter framework. However, the proposed new programme does acknowledge that:

- The NZAF has already done a significant amount of work over its 24 year history to build positive social environments; develop personal skills and strengthen community action.
- The changing HIV epidemic and improvements in HIV treatment mean that people living with HIV will live well and for longer, increasing the potential risk of onward transmission for each positive person – this means the HIV prevention response must have a greater impact than it currently does to reverse the trend of increasing HIV infection rates.
- Information-only approaches (either through health education sessions or resources), although still a necessary

component of a wider programme, on their own have limited impact on behaviour change.

- The increased challenge for HIV prevention as a result of the changing epidemic requires that the NZAF use the resources available in a way that most efficiently and cost-effectively leads to behaviour change, and some activities within current programmes may not be the most effective response.

Improvements in treatment mean that a person may now live with HIV for forty years or more. Rather than being removed from the sexual network upon death at around ten years, HIV will remain present for significantly longer. Condom use throughout an HIV positive person's lifetime must be supported and sustained. This means that NZAF initiatives must be appropriate for those people who are HIV positive. The increase in HIV prevalence increases the risk of infection among risk populations. Put simply, the NZAF must work much harder to reduce the rate of new infection.

Health Information

Many early health promotion approaches to risk behaviour, including HIV prevention approaches were based on an information deficit model. The principal assumption of this model was the belief that 'providing information' about the negative consequences of risks, would effectively deter risk behaviour (e.g. if we tell you unprotected anal intercourse is high risk for HIV infection, you will not have unprotected anal intercourse). For this to be effective, a necessary additional belief was that people behave in a rational manner and that given this new information, they would then alter their behaviour accordingly. It is now clear that both these assumptions are seriously flawed and that the 'information only strategy' by itself is ineffective. (Paglia et al, 1999; Bonell et al, 2000).

While providing information does increase knowledge and awareness, and at times negative attitudes, it does not necessarily have a direct impact on behaviour. Furthermore, providing information about the dangers and risks may even be counterproductive for those who seek adventure, and it may also arouse curiosity in others. Though an information-only approach is not sufficient on its own to affect safe sexual behaviour, it is still recognised that providing facts is a necessary component of a comprehensive HIV prevention programme. Gay men themselves advocate that non-judgemental facts about HIV risk should be provided regularly. (Adams and Neville, 2008).

Events

Building healthy social environments has been a core component of the NZAF HIV prevention programme for many years and the NZAF has been involved in a range of activities including extensive involvement in Pride events in Dunedin, Christchurch, Wellington, Hamilton and Auckland; and involvement in other gay community events.

The new programme proposes a move away from intensive event involvement, and advocates using NZAF sponsorship approaches to support events initiated by others that will carry the core messages of the HIV prevention programme. Additionally, the new programme seeks to ensure that information is integrated into a wider set of activities that carry the core messages (for example, there will not be any brochures or posters that do not fit into a wider programme of work).

Social Marketing

The NZAF HIV prevention response must lead to more men using condoms regularly for anal sex – that means more than the current reality. The proposed new programme has a focus on using more campaign-style media, communications and marketing approaches with a sufficiently large reach to target key populations. New investment is proposed in activities that will build social movements for safe sex; develop cultures of condom use; and market safe sexual behaviour to gay and bisexual men in the environments (both virtual and physical), social networks and media with which they connect. Social marketing approaches provide opportunities for the NZAF to affect more people which leads to behaviour change.

While the NZAF has worked on many campaigns, much of this work has been missing some of the essential ingredients of social marketing. Social marketing uses the techniques of marketing to improve people's health and social conditions. Social marketing is focused on behaviour change, not primarily on improving knowledge and awareness. Significantly, social marketing needs to be sustained over the long term (all previous NZAF campaigns have been three to 12 months in duration). Rather than a focus on the behaviour of the subject, social marketing's starting point is the audience. Social marketing is focused on what the audience takes out, rather than what the organisation wants to put in. Effective social marketing co-ordinates multiple messages across a range of media, and will evaluate and evolve over time. (Bridges and Farlans, 2003).

Results from a range of evaluative research indicate that successful mass media campaigns have a number of common elements, including:

- more effective campaigns use multiple media to promote or challenge a lifestyle norm
- more effective campaigns utilise multiple complementary components and combine and/or link media messages with direct service delivery - e.g. information hotlines and support services and other various prevention efforts
- campaigns should emphasise positive behaviour change rather than negative consequences, and current rewards rather than the avoidance of distant negative consequences
- any anxiety provoking media messages should be accompanied by mechanisms for reducing that anxiety by dealing with it in a productive fashion
- audience segmentation is likely to be more productive if it is based on 'psychographics' (e.g. beliefs, values) rather than on demographics (e.g. social class, gender)
- role models should be carefully selected, as it is essential to be sure that the target group identifies with the models selected and that the role models continue to retain their appeal and credibility
- more effective campaigns avoid fear and moral tactics, and blatant 'hard sells' and should avoid using the organisational logo when possible. (Paglia et al, 1999; Shanahan, 2008; Bonell et al, 2000; Kotler and Lee, 2008).

The goals, outcomes and activities within this plan will all be part of one overall social marketing programme and will begin with the development and launch of a new social marketing brand. Instead of this brand lasting for six or 12 months as previous NZAF campaigns have, the new brand and social marketing programme would last the length of the NZAF HIV Prevention Plan 2009-2014.

The Plan Te Mahere

Outcome Based Planning

The plan has been developed using a programme logic approach, also referred to as outcomes-based planning. A health outcome indicates a change in the health status of an individual or population that is attributable to a planned programme or programmes, regardless of whether a specific individual programme was intended to change that particular health status. The increased use of health outcomes in public health planning reflects a growing understanding that organisational goals alone will not influence actions that will lead to the desired result. Activities that will result in a desirable health outcome for a population must be carried out by many people in a variety of organisations over a number of years.

The sections that follow detail:

- target populations and sub groups
- the overall health goal of the plan
- four target behaviour change goals
- health outcomes for each of the four behaviour change goals
- activities that will be undertaken to achieve each of the outcomes.

The activities have been identified based on an understanding that they will lead to the achievement of the health outcome. The plan suggests combined achievement of the health outcomes under a behaviour change goal will lead to achievement of that goal.

The plan will be evaluated by assessing the impact of the activities on the respective health outcome and overall assessment of behaviour change.



Target Audiences

The primary target population for the NZAF HIV Prevention Plan 2009-14 is gay and bisexual men.

In certain activities, audience segmentation will be used to focus on particular sub-groups that have a significant impact on the behaviour change the programme is seeking to effect. Segmenting the audience based on ethnicity alone is not proposed. The literature states that effective segmenting is based on commonalities of values, beliefs and behaviours as they relate to the behaviour change the programme is seeking.

In the development of the new NZAF HIV Prevention Plan 2009-14, particular groups have been focused on:

- Gay and bisexual men over thirty years of age who have high numbers of sexual partners (more than twenty sexual partners in the last six months)
- Gay and bisexual men who use condoms regularly (around 60%)
- Gay and bisexual men under thirty years of age (the only age cohort with a declining rate of condom use .

Using this segmentation in certain interventions or activities does not imply that other groups will be neglected. For example, there is a focus on men under thirty years of age for some interventions to increase the declining rate of condom use within this group. However, in the majority of interventions, men over thirty years of age will be the principle audience.

The secondary target audience is New Zealand-based African communities.

Behaviours

The four health behaviours that have been identified that will have the greatest impact on the overall health goal of

reducing the incidence of HIV in Aotearoa New Zealand are:

1. increase rates of condom use for anal sex between men (this goal will have the most significant level of resource and activity)
2. increase STI and HIV testing rates for gay and bisexual men
3. increase rates of condom use for first anal sex between men
4. increase rates of condom use within New Zealand-based African communities.

These behaviours form the four goals for this plan.

Goal 1 Whāinga Tuatahi

Increase rates of condom use for anal sex between men

Outcomes

1. *A condom culture among gay and bisexual men is developed and maintained*
2. *Condoms are used in sex on site venues*
3. *A condom culture is supported by those who influence gay and bisexual men*
4. *Condom use among highly sexualised gay and bisexual men is increased*
5. *Regular condom users maintain healthy behaviours*

Unprotected anal intercourse between gay and bisexual men is the single greatest risk behaviour for HIV transmission in New Zealand. By influencing this behaviour the NZAF stands to make the greatest impact on new HIV transmission. Condom use for anal sex significantly reduces onward HIV transmission. (Pinkerton and Abramson, 1997).

With the highest rate of new HIV diagnoses recorded in New Zealand in 2008, the prevalence of HIV in New Zealand is higher than ever before and understanding the nature of gay and bisexual men's sexual networks is essential to understanding patterns of risk behaviours.

The NZAF, in collaboration with the AIDS Epidemiology Group at the University of Otago, conducts biannual behavioural surveillance in the form of the Gay Auckland Periodic Sex Survey (GAPSS) and Gay Online Sex Survey (GOSS). This provides valuable information about the sexual activity, sexuality, sexual knowledge and other demographic factors that must inform the prevention work on which the NZAF embarks. From GAPSS and GOSS we know that:

- in New Zealand, around 60% of sexually active gay and bisexual men use condoms regularly when they have anal sex with casual partners
- 20% report condom use some or most of the time
- 20% report condom use rarely or never
- 66% of men in relationships do not use condoms with their regular partner. (Saxton et al, 2006 .

This data has remained stable since 2002. It is our task to sustain the large portion of men that regularly use condoms by supporting them to maintain healthy behaviours, while encouraging others to increase their condom use. Even a small increase in the groups reporting condom use regularly or often is likely to have a significant impact on onward HIV transmission.

Developing a Condom Culture

Social learning theory (SLT) asserts that people learn from their own experiences and by observing the actions of others and the benefits of those actions (Glanz and Rimer, 2005). SLT was updated by social cognitive theory (SCT) in which behaviour, cognition, and other personal factors, along with environmental influences operate as interacting determinants that influence one another bidirectionally (Peng et al, 2008; Bandura, 1989). Partnering with appropriate organisations and key individuals that have considerable influence on gay and bisexual men, such as sexual health service providers, key GPs with high numbers of gay and bisexual clientele, and health providers and educators (including Māori and Pacific providers) along with mainstream media communications targeting friends and family of gay and bisexual men, is consistent with the findings of evaluative research that have been conducted on social marketing campaigns. The development of a comprehensive 'condom culture' across all population groups in New Zealand would be an enormously effective advance in sexual health for Aotearoa New Zealand. A condom culture is necessitated in part by a trend toward the assimilation of gay and bisexual men into a hetero-normative society which is often reflected in trends in condom use by men in long term relationships.

Men in Relationships

GAPSS and GOSS data (2002, 2004 and 2006) shows that many gay and bisexual men exist within multiple sexual networks.

Men who are in relationships with other men present a challenge for the HIV prevention response for a variety of reasons, including notions of assimilation (among heterosexual relationships), love, monogamy and perhaps toughest of all to address, trust. Approximately 66% of men in relationships report no condom use with their main partner (Saxton et al, 2006; Saxton, 2008).

However, a high number of men who are in relationships with men also report having other concurrent sexual partners. While condom use with casual partners is often high, when that casual partner becomes a more regular partner (called a 'fuckbuddy' in research and colloquially), then condom use with that partner is more likely to decrease. This is complicated by the effects of concurrency, i.e. that fuckbuddy may have other fuckbuddies with whom he has dispensed condom use with too. International and local evidence shows that around half of new HIV infections are among gay and bisexual men who are in relationships. The NZAF HIV prevention response needs to concentrate on increasing sexual health knowledge, behaviour and skills among men who are in relationships with other men. (Saxton, 2008).

Takatāpui and Whānau

Ensuring appropriate HIV prevention interventions for takatāpui and whānau is important. The NZAF HIV Prevention Plan 2009-14 has a commitment to ensuring all interventions are appropriate for takatāpui and whānau and specific staffing skills and experience will be sought for social marketing and community engagement intervention areas. There will also be key targeted initiatives designed to increase condom use with takatāpui:

- www.broonline.co.nz – an online social networking site which models safe sex for takatāpui tane, whānau and 'friends'
- Hui Takatāpui – the NZAF is committed to building community and leadership among takatāpui and a large scale hui which has been a successful intervention to develop this leadership in the past will continue
- Specific engagement, support and resources for providers of health services to takatāpui and whānau
- A targeted resource for whānau of takatāpui, focused on increasing understanding of sexual and gender diversity and sexual health.

Sex on Site Venues

From GAPSS and GOSS it is known that sex on site venue customers are likely to be older (over 45 years) and many will have high numbers of sexual partners. It is interesting to note that this matches the traditional demographic of new diagnoses among gay and bisexual men in New Zealand (aged 30-49). It is essential then that the NZAF HIV prevention response concentrates on increasing the impact that sex on site venues have on condom use. Work has begun on developing a sex on site Venue Accreditation Programme and it is important that this strategy is fully implemented and maintained. Additionally, the venues themselves provide a critical environment in which to promote condom and lube use and NZAF sexual health testing.

Hooking Up Online

Not all highly sexualised men will be reached by partnering with sex on site venues since a significant proportion of men have turned to the internet to find sexual partners. GOSS has shown that the online sample of respondents were more likely to have casual sex, were most likely not to use condoms for anal sex, and were the least likely to have had an HIV test since their last episode of unprotected anal intercourse. Importantly, this group was likely to be the least knowledgeable about HIV information, and the least connected to traditional concepts of gay communities.

The NZAF HIV prevention response must continue and expand online initiatives. These would include NZAF dedicated websites such as www.broonline.co.nz, a social marketing campaign website, nzaf.org.nz, a high presence on social networking sites such as Facebook, Bebo, Twitter, YouTube, and special initiatives on existing and emerging dating/hook-up sites such as NZDating.com.

Community Engagement

Notions and definitions of community are complex and there is no single or correct answer. Discussions and research were carried out that informed the NZAF HIV Prevention Plan 2009-14 about the changes and developments of 'the gay community' and 'gay communities' in Aotearoa New Zealand. While there was no consensus on what these groupings and group identities were, there is agreement that group interactions of gay and bisexual men have changed significantly over the course of the HIV epidemic.

Working effectively with online technology does not replace the importance of face to face community engagement. However, this plan does acknowledge that face to face contact will not reach the majority of gay and bisexual men. Engaging with those most at risk of HIV requires a combination of different forms of community engagement. Traditional 'physical' or face to face engagement will be complemented by a range of virtual engagement interventions, including NZAF websites, social media, mobile technologies and social marketing campaign material.

A new team will ensure effective community engagement in social environments where gay and bisexual men meet. The team will include a focus on effective relationships with Tangata Whenua and Pasifika Communities.

Porn Depicting Unprotected Anal Intercourse

Bareback pornography has recently become more popular. This appears to be a new genre of porn, different from simply being porn without condoms. This is newly produced porn with an edge, since it is made in the face of all safe sex

messages being promoted, and includes an element of wilful riskiness and danger. This is an area that NZAF HIV prevention must understand better, as well as considering ways to counter depictions of unprotected anal intercourse, such as porn that sexualises condom-use, promoting condom-use porn, and using various media to discourage the promotion of bareback porn.

HIV Prevention and Men Living with HIV

The reality for men living with HIV has changed dramatically over the past 20 years. Early in the AIDS epidemic people with HIV died quickly and treatments were ineffective.

The NZAF HIV Prevention Plan 2009-14 acknowledges the importance of people living with HIV to an overall HIV prevention response. In 2003, the US Centres for Disease Control and Prevention included 'persons diagnosed with HIV and their partners' in new high priority prevention efforts. Advanced treatment technology allows positive people to live longer and with improved quality of life they are likely to remain sexually active for longer too. Today life expectancy with successful treatment is thought to be around 40 years from the onset of treatment. Young men living with HIV will not have experienced the early years of the epidemic which saw dramatic illness and death across the gay community.

Most people living with HIV practice safe sex, however research is showing an increase in clusters of people living with HIV in Western nations that are engaging in risky behaviour.

New Zealand behavioural surveillance shows that gay and bisexual men living with HIV are three times more likely to report unprotected sex than HIV negative men.

'Respondents who had been diagnosed with HIV reported the highest rate of any unprotected anal intercourse with a casual partner in the previous six months, compared to those who had tested HIV negative at their most recent test and those who had never tested for HIV. This result remained significant independently of other factors in the multivariate analysis.'

'A difficulty interpreting this arises as it is not known how many of the unprotected episodes occurred with casual sex partners who were also HIV positive, a practice referred to as 'serosorting' or 'positive-positive' sex. New Zealand safe sex guidelines have consistently emphasised the potential personal and public health risks of discarding condoms during positive-positive sex, including HIV super-infection and infection with other (or repeat) STIs, and have not promoted this as an effective safe sex strategy.' (Saxton, 2008:277)

There are also issues specific to men living with HIV (or more prevalent among them) which, if addressed, would be likely to increase condom use for anal sex including:

- superinfection with new strains of HIV
- low viral load and transmission
- lack of disclosure and criminal prosecutions
- heightened rates of acquisition and the effect of STIs on HIV positive men
- sexual dysfunction (heightened by treatments) and condom use
- remaining pro-sexual while living with HIV
- serodiscordant relationships.

The NZAF HIV Prevention Plan 2009-14 ensures a focus on men living with HIV with social marketing interventions, in collaboration with HIV peer support organisations for people living with HIV.

0 ERALL GOAL • Reduce the Incidence of HIV in Aotearoa New Zealand

1

Increase rates of condom use for anal sex between men

Outcome One: A condom culture among gay and bisexual men is developed and maintained

Key Activities

- Media stories countering condom-less porn
- Ensure sex on site venues do not show porn depicting condom-less sex
- Get it On! sponsorship of key events
- Promotion of key research information from GAPSS and GOSS
- Promotion of condom use within mass communications to high sexualised audiences
- Mass communications in gay and mainstream media include realities and complications of sexual networking
- www.broonline.co.nz
- Hui Takatāpui
- Love Life Fonu

Outcome Two: Condoms are used in sex on site venues

Key Activities

- Develop and implement sex on site venue accreditation programme
- Touchscreen technology in sex on site venues
- Promote NZAF sexual health testing

Outcome Three: A condom culture is supported by those that influence gay and bisexual men

Key Activities

- Condoms in taxis project in Auckland, Wellington and Christchurch
- World AIDS Day
- Partnership with appropriate organisations that are highly influential among gay and bisexual men

- Resources for health providers and educators, including Māori and Pacific providers
- Mainstream media communications targeting whānau and friends of gay and bisexual men
- Partnerships with key providers of sexual health services, including the provision of information for gay and bisexual men within sexual health service settings
- Identify and target key GPs with large numbers of registered gay and bisexual male patients

Outcome Four: Condom use among highly sexualised gay and bisexual men is increased

Key Activities

- Targeted online communications
- Develop social movements of highly sexualised men's communities
- Distribute targeted resources (e.g. Party Packs)
- Get it On! sponsorship of key events
- Maintain and build key venue and dance party organiser relationships
- Online dating initiative developed

Outcome Five: Regular condom users maintain healthy behaviours

Key Activities

- Host Get it On! Big Gay Out to encourage community norms of safe sex
- Develop social movements of gay and bisexual men committed to condom use
- Mass communications affirming regular users
- Small and online media about general gay men's sexual health

Goal 2 Whāinga Tuarua

Increase STI and HIV testing rates for gay and bisexual men

Outcomes

1. Social norms within gay and bisexual men's communities support testing
2. Gay and bisexual men are more aware of how to test
3. HIV and STI testing is more available to gay and bisexual men

GAPSS (2002, 2004 and 2006) and GOSS (2006) has shown that many gay and bisexual men do not know their HIV status and that many men have not had an HIV test for many years. Between 2002 and 2006 the percentage of men who had never tested remained consistent at 24%, the same as in the UK, but lower than in Sydney.

Additionally, men living with HIV in New Zealand are more likely to have been diagnosed with an STI in their lifetime (82.6% of men living with HIV have been diagnosed with an STI compared with 39.8% of HIV negative men) and in the past 12 months (13% of men living with HIV compared with 7.6% of HIV negative men). (Saxton, 2008).

STIs present in the body greatly increase the risk of HIV acquisition and transmission, and early detection could lead to fewer new HIV infections. STI testing also provides an opportunity to encourage HIV testing and to discuss a safer approach to an individual's sexual wellbeing and decision making.

The challenge for an effective HIV prevention response is to ensure there is not a contradictory message; condoms work to stop HIV, but everyone should still need to test for HIV regardless.

The NZAF has a clear view that HIV prevention is entirely within an individual's personal control. The NZAF HIV Prevention Plan 2009-14 seeks to ensure appropriate levels of testing while not undermining the central tenet that condoms and lube work to prevent HIV transmission when used correctly and consistently. Consistency of use is a key challenge, and for those that have unprotected anal sex testing is extremely important.

A range of initiatives to increase access, and encourage HIV and STI testing for those who are at risk, will be undertaken in collaboration with sexual health and HIV support services. These initiatives will complement and work alongside the recent expansion of NZAF Positive Health Services into rapid testing for HIV and syphilis in community settings, syphilis testing in sex on site venues and the new, community-based Men's Sexual Health Clinic at the NZAF Burnett Centre in Auckland.

Peer influence on gay and bisexual men should be harnessed. The concept of 'Bring a Mate to Test' will be explored as a part of this activity.

Overall Goal • Reduce the Incidence of HIV in Aotearoa New Zealand

2

Increase STI and HIV testing rates for gay and bisexual men

Outcome One: Social norms within gay and bisexual men's communities support testing

Key Activities

- 'Bring a Mate to Test' and 'Bring Your Team' schemes
- Increase access to testing services via transport and online booking services
- Social movement/peer outreach work at events to encourage testing
- Rewards for testing

Outcome Two: Gay and bisexual men are more aware of how to test

Key Activities

- Small and online media encouraging testing
- Guerrilla marketing
- Text reminders
- Viral marketing (through email, social media and websites)
- Mates Promotion (via whānau, women friends of gay men)
- Takatāpui 'Ask your Aunties' promotion
- Promotion of 0800 number

Outcome Three: HIV and STI testing is more available to gay and bisexual men

Key Activities

- Partnership and shared projects with NZAF Positive Health Services on the delivery and promotion of community-based and venue-based testing services

Goal 3 Whāinga Tuatoru

Increase rates of condom use for first anal sex between men

Outcomes

1. Resiliency among young same sex attracted men is increased

In 2008, there were 12 new homosexually acquired HIV diagnoses in the 20-29 age bracket who were infected in New Zealand. This represents 19% of locally acquired HIV diagnoses among gay and bisexual men, for that year.

GAPSS and GOSS (2002, 2004 and 2006) has found that men who use condoms for anal sex the first time they have anal intercourse are far more likely to continue using condoms for anal sex over the course of their lives. If this is considered alongside the findings from GOSS (2006), which illustrates that around 50% of gay and bisexual men have had anal intercourse by the time they are twenty years old, and around 96% by the time they are thirty, young gay and bisexual men are a key sub-group that must not be neglected.

OVERALL GOAL • Reduce the Incidence of HIV in Aotearoa New Zealand

3

Increase rates of condom use for first anal sex between men

Outcome One: Resiliency among young same sex attracted men is increased

Key Activities

- Partnerships with youth services, sexual health and education providers to increase understanding of, and normalise, anal sex
- Partnership with Rainbow Youth to promote queer youth issues within the framework of OUT THERE!
- www.curious.org.nz Text and Web Service
- Mass communications developed that target tertiary students
- Positive role models for safe sex
- Sponsorship of gay and/or youth events or activities
- Online targeted communications via social media

Goal 4 Whāinga Tuawha

Increase rates of condom use within New Zealand-based African communities

Outcomes

1. Awareness within African communities of HIV risk behaviours is increased
2. Key community leaders are encouraged to influence condom use
3. Sexual health knowledge and skills of African communities is increased
4. Understanding of HIV risk within New Zealand's African communities is developed

In New Zealand, the second population group most affected by HIV are the African communities based in New Zealand. The vast majority of HIV transmission occurred overseas for people living with HIV from African communities.

African people living with HIV in New Zealand are almost all relatively new migrants. The number of refugees living with HIV coming to New Zealand continues to be very small in number.

However, while onward transmission within New Zealand-based African communities remains very low, it is important to keep focused on these communities given the relatively high number of people living with HIV. As of 2008, epidemiology data reports there were 440 people from African communities that had tested HIV positive (223 male and 217 female) living in New Zealand.

Overall Goal • Reduce the Incidence of HIV in Aotearoa New Zealand

4

Increase rates of condom use within New Zealand-based African communities

Outcome One: Awareness within African communities of HIV risk behaviours is increased

Key Activities

- Mass communications in line with NZAF social marketing plans
- Distribute condoms at key and strategic community events
- Sponsorship of, and participation in, key community events
- African prevalence rates distributed among African communities

Outcome Two: Key community leaders are encouraged to influence condom use

Key Activities

- Religious Leaders' Workshops
- Engagement and interaction with key community leaders
- Participation in key community events
- Celebrity modelling of behaviour (e.g. sponsorship of Miss Africa events)
- Youth role models

Outcome Three: Sexual health knowledge and skills of African communities is increased

Key Activities

- Community Educators' Programme
- Dissemination of HIV information to key stakeholders and Community Educators
- Christchurch Women's Project

Outcome Four: Understanding of HIV risk within New Zealand's African communities is developed

Key Activities

- Collaboration with academic institutions on small scale research projects that develop a knowledge base related to HIV risk for people from African communities based in New Zealand.

Programme Evaluation Framework

Aromātai Hātaka

Evaluating a programme of work with as many elements as this plan is complex. A range of evaluation activities has been identified.

Evaluation of this plan will be guided by the programme logic model outlined on page 22. Each evaluation activity will assess the extent to which the four behaviour change goals have been achieved:

1. Increase rates of condom use for anal sex between men.
2. Increase STI and HIV testing rates for gay and bisexual men.
3. Increase rates of condom use for first anal sex between men.
4. Increase rates of condom use within New Zealand-based African communities.

Social marketing evaluation requires work at various levels of intervention. Donovan (2003) suggests that social marketing programmes should measure immediate outcomes in attitudes, beliefs and self-reported behaviour changes.

Evaluation Activities

Four areas of evaluation activity make up the evaluation framework:

Behavioural Surveillance (Behavioural)

GAPSS and GOSS act as a broad outcome evaluation of HIV prevention among gay and bisexual men in Aotearoa New Zealand, providing population level indicators for changes in condom use over time. Behavioural surveillance data for New Zealand-based African communities is lacking. The NZAF will seek to work with potential partners to improve the HIV behavioural surveillance for African communities.

Social Marketing Outcome Evaluation (Outcome)

Population level assessment of behaviour change alongside exposure to messages/branding. A biannual evaluation exercise (in the alternate year to GAPSS and GOSS) will be conducted among gay and bisexual men which will use a methodology that compares the number of exposures to NZAF prevention interventions/resources/messages with the level of desired behaviour (condom use, condom use for first anal sex, HIV and STI testing).

The methodology will be developed in more detail during 2009 and the first outcome evaluation will be completed during Summer 2010-2011.

Impact Evaluation (Impact)

Impact evaluations that measure the intended and unintended effects of specific HIV prevention activities/projects will be completed as one-off evaluation projects over the three to five year duration of this plan. External evaluation consultants and experts may be used. They will include:

- Bro Online
- Sex on Site venue work
- Live Life Fono
- Big Gay Out
- Curious Web and Text Service
- African Youth Workshops
- African Community Educators' Programme.

Practice Evaluation (Practice)

Varcoe (2004) suggests a range of data collection opportunities to assess the impact of social marketing programmes (see table on following page). This will be implemented at the 'practitioner level' within the NZAF staff, ensuring ongoing assessment of our work against the four behaviour change goals of the plan.

Summary

The table below summarises the range of evaluation activities within this framework.

PREVENTION PLAN GOAL	EVALUATION COMPONENT	EVALUATION ACTIVITY	RESULT LEVEL
1 Increase rates of condom use for anal sex between men	Behavioural	GAPSS and GOSS	% increase in condom use.
	Outcome	Social Marketing Outcome Evaluation	Correlation between social marketing programme messaging and this behaviour change goal.
	Impact	Single independent evaluation at one point between 2010-2014 of Bro Online; sex on site venue work; the Love Life Fono and the Big Gay Out	Impact of Love Life Fono on outcome one.
			Impact of range of work with sex on site venues on outcome two. Impact of Bro Online on outcome one. Impact of Big Gay Out on outcome five.
Practice	Web analytics data collection	% increase in online activity or engagement	
2 Increase STI and HIV testing rates for gay and bisexual men	Practice	Testing data collection from NZAF Regional Centres and national HIV testing laboratory data	% increase in STI and HIV testing rates
	Outcome	Social Marketing Outcome Evaluation	Correlation between social marketing programme messaging and this behaviour change goal

PREVENTION PLAN GOAL	EVALUATION COMPONENT	EVALUATION ACTIVITY	RESULT LEVEL
3 Increase rates of condom use for first anal sex between men	Behavioural	GAPSS and GOSS	% increase in condom use for first time anal sex
	Outcome	Social Marketing Outcome Evaluation	Correlation between social marketing programme messaging and this behaviour change goal.
	Impact	Single independent evaluation at one point between 2010-2014 of Curious Web and Text Service	Impact of Curious on outcome one.
	Practice	Web analytics data collection	% increase in online activity or engagement
4 Increase rates of condom use with n New Zealand-based African communities	Behavioural	African behavioural surveillance	% increase in condom use
	Outcome	Social Marketing Outcome Evaluation	Correlation between social marketing programme messaging and this behaviour change goal.
	Impact	Single independent evaluation at one point between 2010-2014 of youth workshops and Community Educators' Programme	Impact of youth workshops on outcome two. Impact of Community Educators' Programme on outcome three.
	Practice	Evaluation of workshops achieving learning outcomes	Increased knowledge of HIV and AIDS reported from workshops (youth and Community Educators' Programme) Evidence of young people acting as role models advocating condom use in their peer groups.

Reference List

Rārangi Pukapuka

- Adams, J. and Neville, S. 2008. *MSM Talk: qualitative research about condom use among MSM in Auckland*. Centre for Social and Health Outcomes Research and Evaluation: Massey University.
- AIDS Epidemiology Group. 2009. *AIDS New Zealand - Issue 63 - March 2009*. Department of Preventive and Social Medicine, University of Otago Medical School: Dunedin.
- AIDS Epidemiology Group. 2008. *AIDS New Zealand - Issue 61 - February 2008*. Department of Preventive and Social Medicine, University of Otago Medical School: Dunedin.
- Auerbach, J. and Coates, T. 2000. 'HIV prevention research: accomplishments and challenges for the third decade of AIDS' in *American Journal of Public Health, Vol 90, Issue 7 1029-1032*. American Public Health Association: USA.
- Bonnell, C., Hickson, F., Hartley, M., Keogh, P., and Weatherburn, P. 2000. *By any means necessary? Reflecting on how HIV prevention interventions work and the changes they bring about*. Sigma Research: London.
- Bourne, A., Dodds, C., Keogh, P., Weatherburn, P., and Hammond, G. 2009. *Relative Safety II: risk and unprotected anal intercourse among gay men with diagnosed HIV*. Sigma Research: London.
- Bridges, T. and Farland, N. 2003. *Social Marketing: Behaviour change marketing in New Zealand*. Senate Communications: Wellington.
- Brown, A., Murphy, G., Rink, G., Clewley, J., Hill, C., Parry, J., Johnson, A., Pillay, D and Gill, O. 2008. 'Implications for HIV testing policy derived from combining data on voluntary confidential testing with viral sequences and serological analyses' in *British Medical Journal 2008;85:4-9*.
- Centre for Disease Control and Prevention. 2003. 'Advancing HIV Prevention : New Strategies for a Changing Epidemic' in *Morbidity and Mortality Weekly Report (MMWR) April 18, 2003 / 52 (15):392-332*. Centre for Disease Control and Prevention: USA.
- Donovan, R. and Henely, N. 2003. *Social Marketing: Principles and Practice*. IP Communications.
- Evans A.R, Wiggins R., and Bolding G, Eiford, J. 2008. 'Characteristics of Gay and Bisexual Men Who Drop Out of a Web Survey of Sexual Behaviour in the UK' in - Springer Science + Business Media, LLC 2008
- Fenton, K. and Bloom, F. 2007. 'STD Prevention with Men Who Have Sex with Men in the United States' in *Behavioural Interventions for Prevention and Control of Sexually Transmitted Diseases 325:353*. Business Media, New York.
- Glanz, K., Rimmer, B., and Biswanath, V. 2008. *Health Behavior and Health Education Theory, Research and Practice*. Jossey Bass: USA.
- Glanz, K., and Rimer, B. 2005. *Theory at a Glance: A guide for Health Promotion Practice*. Department of Health and Human Services, Public Health Service: Washington DC: USA.
- Grierson, J., Thorpe, R., Pitts, M., Hughes, A., Saxton, P., Smith, J., Thomas, M., and Smythe, E. 2008. *HIV Futures New Zealand? Mate Araikore a muri ake nei (Tuarua)*. The Australian Research Centre in Sex, Health and Society, Latrobe University: Melbourne.
- Guy, R., McDonald, A., Barlett, M., Murray, J., Giele, C., Davey, T., Appuhamy, R., Knibbs, P., Coleman, D., Hellard, M., Grulich, A., and Kaldor, J. 2008. 'Characteristics of HIV diagnoses in Australia' in *Sexual Health 5:91-96*. CSIRO Publishing: Australia.
- Hornik, R. 2002. *Public Health Communication: Evidence for Behaviour Change*. Lawrence Erlbaum Associates: Mahwah, NJ.

Holtgrave, D. 2007. 'Evidence Based Efforts to Prevent HIV Infection: An Overview of Current Status and Future Challenges' in *Clinical Infectious Diseases* 45:S293-9. Infectious Diseases Society of America: Arlington, VA.

Kotler, P. and Lee, N. 2009. *Up and Out of Poverty: The Social Marketing Solution*. Wharton School Publishing: USA.

Kotler, P. and Lee, N. 2008. *Social Marketing: Influencing Behaviours for Good*. Sage: USA.

Kotler, P., Roberto, N. and Lee, N. 2002. *Social Marketing: Improving the Quality of Life*. Sage Publications: USA.

Rossen F., Lucassen M., Denny, S. and Robinson, E. 2009. *Youth 07 - The Health and Wellbeing of Secondary School Students in New Zealand*. Auckland Adolescent Health Research Group, University of Auckland: Auckland.

Paglia, A., Room, R. 1999. "Preventing substance-use problems among youth: A literature review and recommendations" in *Journal of Primary Prevention*, vol 20, no.1. pp 3-50

Peng, B. and Schoech, D. 2008. "Grounding Online Prevention Interventions in Theory: Guidelines from a Review of Selected Theories and Research" in *Journal of Technology in Human Services*. Routledge: London.

Pinkerton, S. and Abramson, P. 1997. "Effectiveness of Condoms in Preventing Transmission" in *Social Science and Medicine*, vol 44, no 9. Elsevier Science: UK. pp. 1303-1312.

Plant, A., Montoya, J., Rotblatt, H., Kerndt, P., Mall, K., Pappas, L., Kent, C., and Klausner J. 2008. 'Stop the Sores: The Making and Evaluation of a Successful Social Marketing Campaign' in *Health Promotion Practice* 0:1524839907308376v1. Society for Public Health Education: USA.

Prestage, G., Jin, F., Zablotska, I., Imrie, J., Kaldor, J. and Grulich, A. 2008. 'Trends in HIV prevalence among homosexual and bisexual men in eastern Australian states' in *Sexual Health* 5:103-107. CSIRO Publishing: Australia.

Rosengarten, M., Race, K., and Kippax, S. 2000. *Touch wood, everything will be OK: gay men's understandings of clinical markers in sexual practice*. National Centre in HIV Social Research, Faculty of Arts and Social Sciences, The University of New South Wales: Sydney.

Safren, S., Wingood, G., and Altice, F. 2007. 'Strategies for Primary HIV Prevention That Target Behavioural Change' in *Clinical Infectious Diseases* 2007:45::S300-7. Infectious Diseases Society of America: Arlington, VA.

Sandy, I., Siziya, S., and Fylikesnes, K. 2008. 'Lost opportunities in HIV prevention: programmes miss places where exposures and highest' in *BMC Public Health* 2008, 8:31. BioMed Central.

Saxton, P. 2008. *HIV Epidemiology and Behavioural Surveillance Among Men Who Have Sex with Men in New Zealand*. University of Otago: Dunedin.

Saxton, P., Hughes, T., and Dickson, N. 2006. *GAPSS 2006: Findings from the Gay Auckland Periodic Sex Survey*. New Zealand AIDS Foundation: Auckland.

Shanahan, P., Elliott, B., Dahlgren, N. 2000. *Review of public information campaigns addressing youth risk-taking*. National Youth Affairs Research Scheme: Hobart, Australia.

Sharples, K., Dickson, N., Paul, C., and Skegg, D. 1996. HIV/AIDS in New Zealand: an epidemic in decline? In *AIDS*.10:1273-8.

Spina, A., and Fowler, D. 2007. *Mapping HIV & STI prevention programs and activities targeting gay and other homosexually active men in 2005/06*. Australian Federation of AIDS Organisations: Sydney.

Varcoe, J. Assessing the effectiveness of social marketing. *Unpublished Paper: ESOMAR® -The World Association of Research Professionals*.

Waa, A., Holibar, F., and Spinola, C. 1998. *Programme Evaluation: an introductory guide for health promotion*. University of Auckland: Auckland.



NEW ZEALAND AIDS FOUNDATION
Te Tūāpapa Mate Āraikore o Aotearoa

www.nzaf.org.nz