

UGANDA Global Youth Tobacco Survey (GYTS) Fact Sheet(Ages 13-15)





The Uganda GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Uganda could include in a comprehensive tobacco control program.

The Uganda GYTS was a school-based survey of students in forms S.1, S.2 and S.3 conducted in 2007.

A two-stage cluster sample design was used to produce representative data for Uganda. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 96.1%, the class response rate was 100.0%, the student response rate was 84.5%, and the overall response rate was 81.2%. A total of 2,251 students aged 13-15 participated in the Uganda GYTS.

Prevalence

15.6% of students had ever smoked cigarettes (Boy = 19.2%, Girl = 11.2%)

16.6% currently use any tobacco product (Boy = 17.3%, Girl = 15.3%)

5.5% currently smoke cigarettes (Boy = 6.6%, Girl = 4.0%)

13.9% currently use other tobacco products (Boy = 13.8%, Girl = 13.5%)

6.7% of never smokers are likely to initiate smoking next year

Access and Availability - Current Smokers

32.3% usually smoke at home

18.8% buy cigarettes in a store

Exposure to Secondhand Smoke (SHS)

20.0% live in homes where others smoke in their presence

45.6% are around others who smoke in places outside their home

48.3% think smoking should be banned from public places

79.2% think smoke from others is harmful to them

11.5% have one or more parents who smoke

2.3% have most or all friends who smoke

Cessation - Current Smokers

70.3% want to stop smoking

76.6% tried to stop smoking during the past year

83.4% have ever received help to stop smoking

Media and Advertising

71.2% saw anti-smoking media messages, in the past 30 days

54.7% saw pro-cigarette ads on billboards, in the past 30 days

48.3% saw pro-cigarette ads in newspapers or magazines, in the past 30 days

12.3% have an object with a cigarette brand logo


10.3% were offered free cigarettes by a tobacco company representative


School

70.5% had been taught in class, during the past year, about the dangers of smoking

62.4% had discussed in class, during the past year, reasons why people their age smoke

69.4% had been taught in class, during the past year, the effects of tobacco use





HIGHLIGHTS

- 16.6% of students currently use any form of tobacco; 5.5% currently smoke cigarettes; 13.9% currently use some other form of tobacco.
- SHS exposure is high 1 in 5 students live in homes where others smoke, and close to half the students are exposed to smoke around others outside of the home; 11.5% of students have at least one parent who smokes.
- Eight in 10 students think smoke from others is harmful to them.
- Almost half the students think smoking in public places should be banned.
- Seven in 10 current smokers want to stop smoking.
- One in 8 students has an object with a cigarette brand logo on it.
- Seven in 10 students saw anti-smoking media messages in the past 30 days; over half the students saw pro-cigarette ads on billboards and in newspapers and magazines in the past 30 days.