

Premature mortality from noncommunicable disease



- 10% reduction in premature mortality from injuries & violence, cancer, cardiovascular diseases, diabetes and chronic respiratory diseases by 2020 (set 2017)

Harmful alcohol use



- 10% reduction in harmful use of alcohol by 2020 (set 2017)

Physical inactivity



- 10% reduction in prevalence of insufficient physical activity by 2020 (set 2017)

Sodium intake



- 30% reduction in mean population salt intake by 2020 (set 2017)

Tobacco use



- 30% reduction in prevalence of current tobacco use in persons aged 18+ by 2020 (set 2017)

Raised blood pressure



- 10% reduction in the prevalence of high blood pressure by 2020 (set 2017)

Diabetes and obesity



- 10% reduction in the prevalence of diabetes by 2020 (set 2017)

- 10% reduction in the prevalence of obesity by 2020 (set 2017)

Drug therapy to prevent heart attacks and strokes



- At least 30% of eligible people receive drug therapy and counseling on suicide, mental illness, violence, injuries, harmful use of alcohol, cancer (including glycemic control) to prevent heart attacks and strokes by 2020 (set 2017)

Essential noncommunicable disease medicines and basic technologies to treat major noncommunicable diseases



- 95% availability of the affordable basic technologies and essential medicines, including generics required to treat major non-communicable diseases in public facilities by 2020 (set 2017)