

# Malawi-Lilongwe (Ages 13-15) Global Youth Tobacco Survey (GYTS)



## FACT SHEET . . . . .

The Malawi-Lilongwe GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Malawi-Lilongwe could include in a comprehensive tobacco control program.

The Malawi-Lilongwe GYTS was a school-based survey of students in Standards 7 and 8, and Form 1 conducted in 2000.

A two-stage cluster sample design was used to produce representative data for all of Malawi-Lilongwe. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 100.0%, the class response rate was 100%, the student response rate was 84.0%, and the overall response rate was 84.0%. A total of 1,083 students aged 13-15 years participated in the Malawi-Lilongwe GYTS.

### Prevalence

- 18.9% of students had ever smoked cigarettes (Boys = 28.5%, Girls = 9.8%)
- 16.9% currently use any tobacco product (Boys = 19.1%, Girls = 14.4%)
- 6.1% currently smoke cigarettes (Boys = 9.5%, Girls = 3.1%)
- 12.9% currently use other tobacco products (Boys = 12.8%, Girls = 12.7%)
- 16.9% of never smokers are likely to initiate smoking next year

### Knowledge and Attitudes

- 48.8% think boys and 20.3% think girls who smoke have more friends
- 2.3% think boys and 2.6% think girls who smoke look more attractive

### Access and Availability - Current Smokers

- 30.4% usually smoke at home
- 17.6% buy cigarettes in a store
- 48.7% who bought cigarettes in a store were NOT refused purchase because of their age\*

### Exposure to Secondhand Smoke (SHS)

- 16.0% live in homes where others smoke in their presence
- 35.5% are around others who smoke in places outside their home
- 85.1% think smoking should be banned from public places
- 81.8% think smoke from others is harmful to them
- 8.3% have one or more parents who smoke
- 4.2% have most or all friends who smoke

### Cessation - Current Smokers

- 68.7% want to stop smoking\*
- 88.9% have ever received help to stop smoking

### Media and Advertising

- 85.7% saw anti-smoking media messages, in the past 30 days
- 55.8% saw pro-cigarette ads on billboards, in the past 30 days
- 64.0% saw pro-cigarette ads in newspapers or magazines, in the past 30 days
- 16.9% have an object with a cigarette brand logo
- 14.4% were offered free cigarettes by a tobacco company representative

### School

- 68.9% had been taught in class, during the past year, about the dangers of smoking
- 50.7% had discussed in class, during the past year, reasons why people their age smoke
- 68.2% had been taught in class, during the past year, the effects of tobacco use

\* sample size <35 participants

### Highlights

- More than 1 in 10 students currently use some form of tobacco; 6.1% currently smoke cigarettes; 12.9% currently use some other form of tobacco.
- SHS exposure is moderate – more than 1 in 10 students live in a home where others smoke; more than 3 in 10 are exposed to smoke in public places; 8.3% have one or more parents who smoke.
- More than 8 in 10 students think smoke from others is harmful to them.
- Over 4 in 5 students think smoking should be banned from public places.
- More than two-thirds of smokers want to stop.
- One 1 in 6 students has an object with a cigarette brand logo on it.
- Over 8 in 10 students saw anti-smoking media messages in the past 30 days; more than half of students saw pro-cigarette ads on billboards and almost two-thirds of students saw pro-cigarette ads in newspapers or magazines in the past 30 days.