



The 2013 Cambodia GSHS was a school-based survey of students in grades 7 - 12, which are typically attended by students aged 13 - 17 years. A two-stage cluster sample design was used to produce data representative of all students in grades 7 - 12 in Cambodia. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate.

The Cambodia GSHS measured alcohol use; dietary behaviors; drug use; hygiene; mental health; physical activity; sexual behaviors; tobacco use; and violence and unintentional injury. Students self-reported their responses to each question on a computer scanable answer sheet. The school response rate was 100%, the student response rate was 85%, and the overall response rate was 85%. A total of 3806 students participated in the Cambodia GSHS. Prevalence estimates (percentages) and 95% confidence intervals are presented below.

	Students Aged 13-15 Years			Students Aged 16-17 Years			Students Aged 13-17 Years		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
Alcohol Use									
Among students who ever had a drink of alcohol (other than a few sips), the percentage who had their first drink of alcohol before age 14 years	52.0 (42.4-61.5)	46.2 (36.6-56.0)	*	12.6 (8.2-18.9)	13.8 (8.4-21.9)	10.0 (4.9-19.4)	30.3 (22.8-38.9)	28.4 (21.5-36.6)	34.4 (23.3-47.4)
Percentage of students who drank at least one drink containing alcohol on one or more of the past 30 days	5.4 (4.3-6.9)	7.7 (5.6-10.4)	3.3 (2.2-4.8)	10.5 (8.0-13.7)	14.4 (10.5-19.5)	6.0 (3.9-9.1)	7.3 (6.0-8.8)	10.3 (8.2-12.8)	4.2 (3.1-5.8)
Percentage of students who drank so much alcohol that they were really drunk one or more times during their life	4.5 (3.1-6.4)	6.2 (4.2-9.0)	2.9 (1.8-4.5)	10.5 (7.8-13.9)	15.0 (10.4-21.0)	5.4 (3.9-7.6)	6.7 (5.3-8.4)	9.6 (7.4-12.3)	3.8 (2.8-5.1)
Dietary Behaviors									
Percentage of students who were underweight (<-2SD from median for BMI by age and sex)	14.9 (10.8-20.1)	17.4 (11.1-26.0)	12.6 (10.2-15.6)	13.1 (10.0-17.1)	17.7 (13.0-23.7)	8.0 (5.9-10.7)	14.2 (10.8-18.5)	17.5 (12.4-24.0)	11.0 (8.8-13.6)
Percentage of students who were overweight (>+1SD from median for BMI by age and sex)	3.7 (2.4-5.7)	3.4 (1.9-5.9)	4.0 (2.6-6.1)	3.7 (2.5-5.2)	4.2 (2.4-7.1)	3.1 (2.0-4.8)	3.7 (2.7-5.1)	3.7 (2.4-5.7)	3.7 (2.7-5.0)
Percentage of students who were obese (>+2SD from median for BMI by age and sex)	0.4 (0.3-0.5)	0.5 (0.2-1.2)	0.3 (0.1-1.0)	0.5 (0.2-1.3)	0.8 (0.3-2.4)	0.1 (0.0-0.6)	0.4 (0.3-0.7)	0.6 (0.3-1.3)	0.2 (0.1-0.6)
Percentage of students who usually drank carbonated soft drinks one or more times per day during the past 30 days	45.6 (41.0-50.3)	45.0 (37.7-52.5)	46.4 (41.7-51.1)	41.8 (37.0-46.7)	41.3 (35.9-47.0)	42.3 (36.3-48.5)	44.2 (40.0-48.5)	43.6 (38.4-48.9)	44.9 (40.5-49.4)



	Students Aged 13-15 Years			Students Aged 16-17 Years			Students Aged 13-17 Years		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
Drug Use									
Percentage of students who used marijuana one or more times during their life	1.2 (0.6-2.5)	1.2 (0.6-2.7)	1.2 (0.5-2.8)	0.8 (0.3-1.9)	1.3 (0.5-3.4)	0.2 (0.0-1.8)	1.1 (0.6-2.0)	1.3 (0.7-2.4)	0.9 (0.4-2.0)
Hygiene									
Percentage of students who usually cleaned or brushed their teeth less than one time per day during the past 30 days	4.9 (3.7-6.6)	5.4 (4.0-7.4)	4.3 (2.5-7.3)	3.8 (2.4-6.2)	5.2 (3.2-8.4)	2.3 (1.0-5.2)	4.5 (3.3-6.2)	5.3 (3.9-7.3)	3.6 (2.3-5.7)
Percentage of students who never or rarely washed their hands after using the toilet or latrine during the past 30 days	3.1 (2.1-4.3)	3.6 (2.4-5.3)	2.5 (1.6-4.0)	2.6 (1.1-6.1)	3.7 (1.3-10.5)	1.3 (0.5-3.2)	2.9 (2.1-3.9)	3.7 (2.5-5.3)	2.1 (1.4-3.2)
Mental Health									
Percentage of students who ever seriously considered attempting suicide during the past 12 months	5.5 (4.2-7.1)	4.2 (2.8-6.4)	6.7 (5.0-8.9)	7.6 (6.4-8.9)	7.0 (5.1-9.4)	8.2 (5.8-11.6)	6.2 (5.4-7.2)	5.3 (3.9-7.1)	7.2 (6.1-8.5)
Percentage of students who actually attempted suicide one or more times during the past 12 months	7.4 (5.9-9.3)	6.4 (4.4-9.2)	8.4 (6.5-10.9)	6.3 (5.0-7.9)	5.0 (3.4-7.3)	7.7 (5.9-10.0)	7.0 (5.8-8.4)	5.9 (4.2-8.1)	8.2 (6.8-9.9)
Percentage of students who had no close friends	5.7 (4.4-7.3)	6.7 (5.0-8.8)	4.7 (3.2-7.1)	4.3 (3.1-5.9)	5.3 (3.2-8.6)	3.2 (1.9-5.5)	5.2 (4.2-6.3)	6.1 (4.5-8.2)	4.2 (3.1-5.8)
Physical Activity									
Percentage of students who were physically active for a total of at least 60 minutes per day on five or more days during the past seven days	9.3 (7.6-11.2)	10.6 (8.6-13.0)	8.0 (5.6-11.4)	10.8 (8.3-13.9)	12.6 (9.5-16.6)	8.7 (5.1-14.5)	9.8 (8.3-11.7)	11.4 (9.7-13.4)	8.3 (5.9-11.4)
Percentage of students who went to physical education (PE) class on three or more days each week during this school year	11.3 (8.7-14.6)	12.8 (9.7-16.6)	10.0 (6.9-14.2)	10.3 (7.9-13.4)	12.2 (8.5-17.2)	8.2 (5.5-12.0)	10.9 (9.1-13.0)	12.5 (10.4-15.1)	9.3 (6.8-12.8)
Percentage of students who spent three or more hours per day during a typical or usual day doing sitting activities	10.5 (7.9-13.7)	11.0 (8.3-14.4)	10.0 (6.8-14.5)	14.1 (11.5-17.1)	14.3 (10.5-19.2)	13.8 (10.6-17.8)	11.8 (9.6-14.5)	12.3 (9.8-15.3)	11.4 (8.8-14.6)



	Students Aged 13-15 Years			Students Aged 16-17 Years			Students Aged 13-17 Years		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
Sexual Behaviors									
Percentage of students who ever had sexual intercourse	13.0 (10.8-15.6)	15.0 (11.6-19.3)	11.2 (8.7-14.2)	10.1 (8.3-12.2)	13.6 (10.5-17.4)	6.3 (4.5-8.7)	11.9 (10.1-14.0)	14.5 (11.8-17.6)	9.4 (7.6-11.7)
Tobacco Use									
Among students who ever smoked cigarettes, the percentage who first tried a cigarette before age 14 years	*	*	*	*	*	*	57.7 (44.5-69.9)	*	*
Percentage of students who smoked cigarettes on one or more days during the past 30 days	2.4 (1.5-3.7)	2.9 (1.7-4.9)	1.9 (1.1-3.2)	2.5 (1.5-4.1)	4.2 (2.4-7.1)	0.6 (0.1-3.1)	2.4 (1.7-3.4)	3.4 (2.3-5.0)	1.4 (0.8-2.6)
Percentage of students who reported people smoked in their presence on one or more days during the past seven days	46.4 (43.7-49.0)	49.5 (45.5-53.6)	43.3 (39.7-47.0)	53.4 (49.5-57.2)	56.7 (50.9-62.2)	49.7 (44.5-54.9)	49.0 (46.4-51.6)	52.3 (48.4-56.2)	45.6 (42.7-48.5)
Violence and Unintentional Injury									
Percentage of students who were in a physical fight one or more times during the past 12 months	13.8 (11.7-16.2)	15.3 (12.1-19.2)	12.2 (9.4-15.6)	8.0 (6.1-10.6)	8.2 (6.2-10.7)	7.9 (5.0-12.3)	11.6 (10.2-13.3)	12.5 (10.2-15.2)	10.7 (8.4-13.5)
Percentage of students who were seriously injured one or more times during the past 12 months	20.2 (16.9-24.0)	21.8 (16.4-28.3)	18.7 (14.4-23.9)	19.2 (15.9-23.1)	22.3 (17.3-28.1)	15.7 (11.8-20.7)	19.9 (17.0-23.0)	22.0 (17.6-27.1)	17.7 (14.2-21.7)
Percentage of students who were bullied on one or more days during the past 30 days	22.4 (18.5-26.8)	22.5 (19.2-26.0)	22.2 (16.1-29.9)	22.3 (17.4-28.2)	21.6 (15.9-28.8)	23.2 (17.1-30.6)	22.4 (18.6-26.6)	22.1 (19.2-25.4)	22.5 (17.1-29.1)

*Indicates data were not available.

For additional information about the Cambodia GSHS, please contact:

Dr. Chher Tepirou
Ministry of Health, Cambodia
tepirou@yahoo.com

For general information about the GSHS, please visit:

<http://www.who.int/chp/gshs/en/>
or
<http://www.cdc.gov/gshs/>