



Congo (Brazzaville) STEPS Survey 2004

Fact Sheet

The STEPS survey of chronic disease risk factors in Congo was carried out in May 2004. Congo carried out Step 1, Step 2 and Step 3. Socio demographic and behavioural information was collected in Step 1. Physical measurements such as height, weight and blood pressure were collected in Step 2. Biochemical measurements were collected to assess blood glucose levels in Step 3 only among participants with risk factors for diabetes (older than 40 years, obese, having raised blood pressure, or family history of hypertension or diabetes). The STEPS survey in Congo was a population-based survey of adults aged 25-64. A cluster sample design was used to produce representative data for that age range in Congo. A total of 2,030 adults participated in the Congo STEPS survey.

Results for adults aged 25-64 years (incl. 95% CI)	Both Sexes	Males	Females
Step 1 Tobacco Use			
Percentage who currently smoke tobacco	11.1%	20.4%	1.7%
Percentage who currently smoke tobacco daily	8.2%	15.5%	0.9%
<i>For those who smoke tobacco daily</i>			
Average age started smoking (years)	20.8	20.4	*
Percentage smoking manufactured cigarettes	88.8%	89.5%	*
Mean number of manufactured cigarettes smoked per day	3.4	3.5	*
Step 1 Alcohol Consumption			
Percentage of abstainers (who did not drink alcohol in the last year)	40.1%	31.1%	49.2%
Percentage of current drinkers (who drank alcohol in the past 30 days)	--	--	--
<i>Among past 12 month drinkers</i>			
Percentage who drank alcohol on 4 or more days in the last week	8.3%	10.7%	3.8%
Percentage who binge drink (men who had 5 or more / women who had 4 or more drinks on any day in the last week)	--	6.8%	7.1%
Step 1 Fruit and Vegetable Consumption (in a typical week)			
Mean number of days fruit consumed	2.8	2.8	2.7
Mean number of servings of fruit consumed on average per day	--	--	--
Mean number of days vegetables consumed	3.6	3.5	3.8
Mean number of servings of vegetables consumed on average per day	--	--	--
Percentage who ate less than 5 combined servings of fruit and/or vegetables on average per day	--	--	--
Step 1 Physical Activity			
Percentage with low levels of activity (defined as <600 MET-minutes/week)**	--	--	--
Percentage with high levels of activity (defined as ≥ 3000 MET-minutes/week)**	--	--	--
Median time spent in physical activity per day (minutes)	--	--	--
Percentage not engaging in vigorous physical activity	--	--	--

** For complete definitions of low and high levels of physical activity, other conditions are specified in the GPAQ Analysis Guide, available at: <http://www.who.int/chp/steps/GPAQ/en/index.html>



Congo (Brazzaville) STEPS Survey 2004

Fact Sheet

Results for adults aged 25-64 years (incl. 95% CI)	Both Sexes	Males	Females
Step 2 Physical Measurements			
Mean body mass index - BMI (kg/m ²)	23.2	22.0	24.6
Percentage who are overweight (BMI ≥25 kg/m ²)	27.3%	18.0%	37.1%
Percentage who are obese (BMI ≥30 kg/m ²)	8.6%	2.5%	15.0%
Average waist circumference (cm)	-	81.6	85.3
Mean systolic blood pressure - SBP (mmHg), including those currently on medication for raised BP	127.7	129.5	125.8
Mean diastolic blood pressure - DBP (mmHg), including those currently on medication for raised BP	83.3	83.8	82.8
Percentage with raised BP (SBP ≥140 and/or DBP ≥90 mmHg or currently on medication for raised BP)	33.3%	34.1%	32.5%
Percentage with raised BP (SBP ≥160 and/or DBP ≥100 mmHg or currently on medication for raised BP)	17.3%	16.6%	18.0%
Step 3 Biochemical Measurement			
Mean fasting blood glucose [mmol/L], excluding those currently on medication for raised blood glucose	4.5	4.6	4.4
Percentage with impaired fasting glycaemia as defined below <ul style="list-style-type: none"> • plasma venous value ≥6.1 mmol/L (110mg/dl) and <7.0 mmol/L (126 mg/dl) • capillary whole blood value ≥5.6 mmol/L (100mg/dl) and <6.1 mmol/L (110 mg/dl) 	3.6%	5.0%	2.6%
Percentage with raised blood glucose as defined below or currently on medication for raised blood glucose <ul style="list-style-type: none"> • plasma venous value ≥7.0 mmol/L or ≥126 mg/dl • capillary whole blood value ≥6.1 mmol/L or ≥110 mg/dl 	20.8%	20.0%	21.4%
Mean total blood cholesterol [choose accordingly: mmol/L or mg/dl]	--	--	--
Percentage with raised total cholesterol (≥5.0 mmol/L or ≥190 mg/dl)	--	--	--
Percentage with raised total cholesterol (≥6.2 mmol/L or ≥240 mg/dl)	--	--	--
Summary of combined risk factors			
<ul style="list-style-type: none"> <li style="width: 50%;">• current daily smokers <li style="width: 50%;">• overweight (BMI ≥ 25 kg/m²) <li style="width: 50%;">• less than 5 servings of fruits and/or vegetables on average per day <li style="width: 50%;">• raised BP (SBP ≥ 140 and/or DBP ≥ 90 mmHg or currently on medication for raised BP) <li style="width: 50%;">• Low level of activity (<600 MET -minutes) 			
Percentage with none of the risk factors included above	--	--	--
Percentage with at least three of the risk factors included above, aged 25 to 44 years old	--	--	--
Percentage with at least three of the risk factors included above, aged 45 to 64 years old	--	--	--
Percentage with at least three of the risk factors included above, aged 25 to 64 years old	--	--	--

* Indicates sample size less than 50.

**For additional information, please contact:
STEPS country focal point Prof. Gisèle Kimbally-Kaky
MOHkimbally2000@yahoo.fr**