

## APPENDICES: FACT SHEET - STEPS Survey 2014

# Uganda STEPS Survey 2014 Fact Sheet

The STEPS survey of non-communicable disease (NCD) risk factors in Uganda was carried out from April, 2014 through June, 2014. Uganda carried out Step 1, Step 2 and Step 3. Socio demographic and behavioral information was collected in Step 1. Physical measurements such as height, weight and blood pressure were collected in Step 2. Biochemical measurements were collected to assess blood glucose and cholesterol levels in Step 3. The survey was a population-based survey of adults aged 18-69. A Multistage sample design was used to produce representative data for that age range in Uganda. A total of 3987 adults participated in the survey. The response rate was 99.0% for STEPs 1 & 2 and 92.2% for all STEPs 1,2 and 3. A repeat survey is planned for 2024 funds permitting.

Results for adults aged 18-69 years (incl. 95% CI)	Both Sexes	Males	Females
<b>Step 1 Tobacco Use</b>			
Percentage who currently smoke tobacco	9.6 (8.1-11.1)	16.8 (14.2-19.5)	2.9 (1.8-4.0)
Percentage who currently smoke tobacco daily	8.3 (6.9-9.8)	14.5 (12.0-17.1)	2.6 (1.5-3.7)
<i>For those who smoke tobacco daily</i>			
Average age started smoking (years)	22.1 (20.9-23.3)	22.2 (20.8-23.6)	21.5 (18.1-24.8)
Percentage of daily smokers smoking manufactured cigarettes	61.5 (53.1-70.0)	68.8 (59.5-78.1)	24.5 (11.3-37.6)
Mean number of manufactured cigarettes smoked per day (by smokers of manufactured cigarettes)	2.7 (2.1-3.2)	3.0 (2.4-3.6)	0.9 (0.2-1.7)
<b>Step 1 Alcohol Consumption</b>			
Percentage who are lifetime abstainers	51.8 (49.3-54.3)	40.4 (36.9-43.8)	62.4 (59.2-65.5)
Percentage who are past 12 month abstainers	12.1 (10.8-13.4)	12.5 (10.4-14.5)	11.8 (10.2-13.4)
Percentage who currently drink (drank alcohol in the past 30 days)	28.5 (26.2-30.8)	40.1 (36.5-43.6)	17.9 (15.3-20.5)
Percentage who engage in heavy episodic drinking (6 or more drinks on any occasion in the past 30 days)	16.7 (14.9-18.5)	26.2 (23.1-29.4)	7.9 (6.3-9.6)
<b>Step 1 Fruit and Vegetable Consumption (in a typical week)</b>			
Mean number of days fruit consumed	2.9 (2.7-3.0)	2.8 (2.6-3.0)	2.9 (2.8-0.0)
Mean number of servings of fruit consumed on average per day	1.4 (1.2-1.5)	1.3 (1.1-1.5)	1.4 (1.2-1.6)
Mean number of days vegetables consumed	3.5 (3.4-3.7)	3.2 (3.0-3.4)	3.8 (3.6-4.0)
Mean number of servings of vegetables consumed on average per day	1.3 (1.2-1.4)	1.2 (1.1-1.3)	1.4 (1.3-1.5)
Percentage who ate less than 5 servings of fruit and/or vegetables on average per day	87.8 (85.9-89.8)	88.4 (85.9-90.8)	87.3 (85.2-89.5)
<b>Step 1 Physical Activity</b>			
Percentage with insufficient physical activity (defined as < 150 minutes of moderate-intensity activity per week, or equivalent)*	4.3 (3.4-5.3)	3.7 (2.1-5.4)	4.9 (3.8-6.0)
Median time spent in physical activity on average per day (minutes) (presented with inter-quartile range)	308.6 (171.4-471.4)	325.7 (184.3-497.1)	285.0 (155.7-441.4)
Percentage not engaging in vigorous activity	49.6 (46.5-52.7)	40.1 (36.3-43.9)	58.4 (54.7-62.1)

<b>Step 1 Cervical Cancer Screening</b>			
Percentage of women aged 30-49 years who have ever had a screening test for cervical cancer			9.9 (6.9-12.8)
<b>Step 2 Physical Measurements</b>			
Mean body mass index - BMI (kg/m <sup>2</sup> )	22.6 (22.4-22.8)	21.7 (21.5-22.0)	23.4 (23.1-23.7)
Percentage who are overweight (BMI ≥ 25 kg/m <sup>2</sup> )	19.1 (17.2-21.0)	11.3 (9.3-13.3)	27.1 (24.2-29.9)
Percentage who are obese (BMI ≥ 30 kg/m <sup>2</sup> )	4.6 (3.7-5.6)	1.8 (1.0-2.6)	7.5 (6.0-9.1)
Average waist circumference (cm)		77.1 (76.6-77.6)	80.3 (79.5-81.0)
Mean systolic blood pressure - SBP (mmHg), including those currently on medication for raised BP	124.5 (123.6-125.3)	126.6 (125.5-127.7)	122.5 (121.5-123.5)
Mean diastolic blood pressure - DBP (mmHg), including those currently on medication for raised BP	80.6 (79.9-81.2)	80.2 (79.3-81.2)	80.9 (80.2-81.6)
Percentage with raised BP (SBP ≥ 140 and/or DBP ≥ 90 mmHg or currently on medication for raised BP)	24.3 (22.3-26.4)	25.8 (22.8-28.9)	22.9 (20.5-25.3)
Percentage with raised BP (SBP ≥ 140 and/or DBP ≥ 90 mmHg) who are not currently on medication for raised BP	93.5 (91.8-95.3)	96.9 (95.4-98.4)	90.1 (87.0-93.2)
<b>Step 3 Biochemical Measurement</b>			
Mean fasting blood glucose, including those currently on medication for raised blood glucose (mmol/l)	3.9 (3.8-4.0)	3.8 (3.7-3.9)	3.9 (3.9-4.0)
Percentage with impaired fasting glycaemia as defined below •capillary whole blood value ≥6.1 mmol/L (110 mg/dl) and <7.0 mmol/L (126 mg/dl)	2.0 (1.5 – 2.5)	2.0 (1.2 – 2.8)	1.9 (1.2 – 2.6)
Percentage with raised fasting blood glucose as defined below or currently on medication for raised blood glucose •capillary whole blood value ≥ 7.0 mmol/L (126 mg/dl)	1.3 (0.7 – 1.9)	1.7 (0.6 – 2.8)	1.0 (0.5 – 1.5)
Mean total blood cholesterol, including those currently on medication for raised cholesterol (mmol/l)	3.4 (3.4-3.5)	3.3 (3.2-3.3)	3.6 (3.5-3.7)
Percentage with raised total cholesterol (≥ 5.0 mmol/L or ≥ 190 mg/dl or currently on medication for raised cholesterol)	6.7 (5.6-7.8)	4.4 (3.1-5.6)	8.9 (7.2-10.5)
Percentage with low HDL Cholesterol (males ≤ 1.03 mmol/l and females ≤ 1.29 mmol/l)		59.9 (56.1 - 63.6)	68.3 (65.4 - 71.2)
<b>Cardiovascular disease (CVD) risk</b>			
Percentage aged 40-69 years with a 10-year CVD risk ≥ 30%, or with existing CVD**	10.5 (8.1-12.9)	6.9 (4.1-9.6)	14.2 (10.6-17.8)
<b>Summary of combined risk factors</b>			
<ul style="list-style-type: none"> <li>• current daily smokers</li> <li>• less than 5 servings of fruits &amp; vegetables per day</li> <li>• insufficient physical activity</li> <li>• overweight (BMI ≥ 25 kg/m<sup>2</sup>)</li> <li>• raised BP (SBP ≥ 140 and/or DBP ≥ 90 mmHg or currently on medication for raised BP)</li> </ul>			
Percentage with none of the above risk factors	6.6 (5.2-8.1)	6.0 (4.2-7.8)	7.3 (5.6-9.0)
Percentage with three or more of the above risk factors, aged 18 to 44 years	6.6 (5.3-7.8)	5.8 (4.2-7.3)	7.4 (5.6-9.2)
Percentage with three or more of the above risk factors, aged 45 to 69 years	20.9 (16.1-25.6)	18.6 (11.1-26.2)	23.0 (17.6-28.3)
Percentage with three or more of the above risk factors, aged 18 to 69 years	9.8 (8.1-11.4)	8.5 (6.3-10.7)	11.0 (9.0-13.0)

\*\* A 10-year CVD risk of ≥30% is defined according to age, sex, blood pressure, smoking status (current smokers OR those who quit smoking less than 1 year before the assessment), total cholesterol, and diabetes (previously diagnosed OR a fasting plasma glucose concentration >7.0 mmol/l (126 mg/dl)).

**For additional information, please contact:**  
**STEPS country focal point STEPS country focal point**  
**[Dr. Gerald Mutungi, [gnmutungi@yahoo.com](mailto:gnmutungi@yahoo.com)]**  
[www.health.go.ug](http://www.health.go.ug)