

Zimbabwe (Manicaland)

2003 Fact Sheet



The **2003 Zimbabwe (Manicaland) GSHS** measured alcohol and other drug use; sexual behaviours that contribute to HIV infections, other STIs, and unintended pregnancies; unintentional injuries and violence; hygiene; physical activity; tobacco use; mental health; and protective factors.

The 2003 Zimbabwe (Manicaland) GSHS was a school-based survey of students in forms 1, 2, and 3. A two-stage cluster sample design was used to produce data representative of all students in forms 1, 2, and 3 in Zimbabwe (Manicaland). At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 100%, the student response rate was 85%, and the overall response rate was 85%. A total of 1,864 students participated in the Zimbabwe (Manicaland) GSHS. Students self-reported their responses to each question on a computer scannable answer sheet. For comparison purposes, only students aged 13-15 years are included in the analyses for this fact sheet.

Results for students aged 13-15 years	Total	Boys	Girls
Alcohol and Other Drug Use			
Percentage of students who had at least one drink containing alcohol on one or more days during the past 30 days	17.0 ± 4.0	20.9 ± 5.6	13.2 ± 4.4
Percentage of students who drank so much alcohol that they were really drunk one or more times during their life	21.3 ± 3.3	25.4 ± 4.5	17.3 ± 5.1
Percentage of students who had a hang-over, felt sick, got into trouble with family or friends, missed school, or got into fights one or more times as a result of drinking alcohol during their life	24.2 ± 6.3	26.3 ± 6.5	22.1 ± 7.6
Percentage of students who used drugs one or more times during their life	15.1 ± 4.8	18.9 ± 6.1	11.3 ± 4.9
Hygiene			
Percentage of students who never or rarely washed their hands before eating during the past 30 days	5.4 ± 2.1	4.7 ± 1.4	6.1 ± 3.2
Percentage of students who never or rarely washed their hands after using the toilet or latrine during the past 30 days	10.4 ± 3.0	12.1 ± 4.5	8.8 ± 2.9
Mental Health			
Percentage of students who felt lonely most of the time or always during the past 12 months	21.3 ± 4.9	22.8 ± 4.2	19.9 ± 6.2
Percentage of students who ever seriously considered attempting suicide during the past 12 months	32.7 ± 6.4	32.8 ± 7.6	32.6 ± 6.2
Percentage of students who have no close friends	13.9 ± 2.0	13.2 ± 3.2	14.5 ± 3.1
Physical Activity			
Percentage of students who were physically active for a total of at least 60 minutes per day on all 7 days during the past 7 days	11.7 ± 3.3	13.0 ± 4.5	10.5 ± 3.4
Percentage of students who spent three or more hours per day during a typical or usual day sitting and watching television, playing computer games, talking with friends, or doing other sitting activities	40.4 ± 4.8	38.9 ± 4.3	42.0 ± 6.8
Protective Factors			
Percentage of students who missed classes or school without permission on one or more days during the past 30 days	37.6 ± 6.6	39.4 ± 7.4	35.9 ± 7.5
Percentage of students who reported that most of the students in their school were never or rarely kind and helpful during the past 30 days	33.1 ± 6.5	37.4 ± 7.6	28.9 ± 6.1
Percentage of students whose parents or guardians never or rarely really knew what they were doing with their free time during the past 30 days	35.5 ± 3.9	37.1 ± 5.6	33.9 ± 5.4

Zimbabwe (Manicaland)

2003 Fact Sheet



Results for students aged 13-15 years	Total	Boys	Girls
Sexual Behaviours that contribute to HIV Infections, Other STIs, and Unintended Pregnancies			
Percentage of students who have ever had sexual intercourse	18.1 ± 4.9	28.0 ± 7.8	9.6 ± 4.1
Percentage of students who ever had sexual intercourse with two or more people during their life	14.9 ± 6.0	22.2 ± 10.0	8.6 ± 3.6
Among student who ever had sexual intercourse, the percentage who used a condom the last time they had sexual intercourse	52.0 ± 8.0	*	*
Tobacco Use			
Percentage of students who smoked cigarettes on one or more days during the past 30 days	10.2 ± 2.3	11.8 ± 3.7	8.6 ± 3.3
Percentage of students who used any tobacco products other than cigarettes on one or more days during the past 30 days	15.8 ± 4.8	17.2 ± 4.3	14.5 ± 6.5
Percentage of students who reported people smoking in their presence on one or more days during the past 7 days	46.2 ± 6.9	47.7 ± 7.8	44.6 ± 7.5
Unintentional Injuries and Violence			
Percentage of students who were in a physical fight one or more times during the past 12 months	47.3 ± 6.2	49.2 ± 7.1	45.5 ± 6.7
Percentage of students who were seriously injured one or more times during the past 12 months	73.2 ± 6.2	74.2 ± 5.9	72.2 ± 8.7
Percentage of students who were bullied on one or more days during the past 30 days	67.2 ± 4.6	65.9 ± 4.0	68.6 ± 6.4

* Indicates less than 100 students

For additional information, please contact:

Dr. Edwin Sithole, Medical Officer
Zimbabwe Assoc. of Church Hospitals, Harare, Zimbabwe, esithole@healthnet.org.zw