



# Algeria STEPS Survey 2016-2017

## Fact Sheet

The STEPS survey on risk factors for non-communicable diseases (NCDs) in Algeria was carried out from November 14, 2016 to May 26, 2017. performed steps 1, 2 and 3. Sociodemographic and behavioral information was collected in step 1. Physical measurements such as height, weight and blood pressure were collected in step 2. Biochemical measurements were collected to assess blood sugar and cholesterol levels in step 3 and Hb1c. The survey was a population-based survey. adults aged 18 to 69. A 3-stage cluster sampling design was used to produce representative data for this age group in Algeria. A total of 6989 adults participated in the survey. The overall response rate was 93.8%. A new investigation is planned for 2022 if funds allow.

| Results for adults aged 18-69 years (incl. 95% CI)   | Both Sexes           | Males                 | Females              |
|--|----------------------|-----------------------|----------------------|
| <b>Step 1 Tobacco Use</b>  |                      |                       |                      |
| Percentage who currently smoke tobacco   | 16.5<br>(15.3-17.6)  | 32.2<br>(30.2-34.3)   | 0.4<br>(0.2-0.7)     |
| Percentage who currently smoke tobacco daily   | 14.3<br>(13.2-15.4)  | 28.1<br>(26.1-30.0)   | 0.3<br>(0.1 – 0.5)   |
| <i>For those who smoke tobacco daily</i>   |                      |                       |                      |
| Average age started smoking (years)  | 17.6<br>(17.3-17.9)  | 17.6<br>(17.3-17.9)   | 18.9<br>(16.2-21.6)  |
| Percentage of daily smokers smoking manufactured cigarettes  | 97.6<br>(96.1-99.0)  | 97.6<br>(96.2-99.1)   | 93.5<br>(79.7-100.0) |
| Mean number of manufactured cigarettes smoked per day (by smokers of manufactured cigarettes)                                    | 14.9<br>(14.2-15.7)  | 14.9<br>(14.2-15.7)   | 15.3<br>(7.6-23.0)   |
| <b>Step 1 Alcohol Consumption</b>  |                      |                       |                      |
| Percentage who are lifetime abstainers   | 91.1<br>(89.9-92.2)  | 82.6<br>(80.3-84.8)   | 99.7<br>(99.5-99.9)  |
| Percentage who are past 12-month abstainers  | 5.6<br>(4.8-6.4)     | 10.9<br>(9.4-12.5)    | 0.2<br>(0 – 0.3)     |
| Percentage who currently drink (drank alcohol in the past 30 days)   | 2.1<br>(1.5-2.7)     | 4.1<br>(3.0-5.2)      | 0.1<br>(0-0.1)       |
| Percentage who engage in heavy episodic drinking (6 or more drinks on any occasion in the past 30 days)                          | 1.3<br>(0.9-1.6)     | 2.5<br>(1.8-3.2)      | 0.0<br>(0-0.1)       |
| <b>Step 1 Diet</b>   |                      |                       |                      |
| Mean number of days fruit consumed in a typical week   | 3.3<br>(3.2-3.4)     | 3.4<br>(3.2-3.5)      | 3.3<br>(3.2-3.4)     |
| Mean number of servings of fruit consumed on average per day   | 0.9<br>(0.9-1.0)     | 1.0<br>(0.9-1.0)      | 0.9<br>(0.8- 1.0)    |
| Mean number of days vegetables consumed in a typical week  | 5.5<br>(5.4-5.6)     | 5.5<br>(5.4-5.6)      | 5.5<br>(5.4-5.6)     |
| Mean number of servings of vegetables consumed on average per day  | 2.1<br>(1.9-2.3)     | 2.2<br>(1.9-2.4)      | 2.1<br>(1.9-2.3)     |
| Percentage who ate less than 5 servings of fruit and/or vegetables on average per day  | 85.3<br>(82.9-87.7)  | 84.8<br>(82.0-87.6)   | 85.9<br>(83.4-88.4)  |
| Percentage who always or often add salt or salty sauce to their food before eating or as they are eating                         | 20.2<br>(18.4-22.0)  | 20.0<br>(17.6-22.3)   | 20.5<br>(18.5-22.5)  |
| Percentage who always or often eat processed foods high in salt  | 17.0<br>(15.3-18.7)  | 19.1<br>(16.8-21.3)   | 15.0<br>(13.2-16.7)  |
| <b>Step 1 Physical Activity</b>  |                      |                       |                      |
| Percentage with insufficient physical activity (defined as <150 minutes of moderate-intensity activity per week, or equivalent)* | 23.7<br>(21.7-25.7)  | 13.7<br>(12.0-15.5)   | 33.7<br>(30.7-36.7)  |
| Median time spent in physical activity on average per day (minutes) (presented with inter-quartile range)                        | 79.3<br>(21.4-197.1) | 120.0<br>(45.7-274.3) | 42.9<br>(8.6-128.6)  |
| Percentage not engaging in vigorous activity   | 69.2<br>(67.1-71.2)  | 54.5<br>(51.9-57.1)   | 83.8<br>(81.4-86.2)  |

\* For complete definitions of insufficient physical activity, refer to theGPAQ Analysis Guide (<http://www.who.int/chp/steps/GPAQ/en/index.html>) or to the WHO Global recommendations on physical activity for health ([http://www.who.int/dietphysicalactivity/factsheet\\_recommendations/en/index.html](http://www.who.int/dietphysicalactivity/factsheet_recommendations/en/index.html))



# Algeria STEPS Survey 2016-2017

## Fact Sheet

| Results for adults aged 18-69 years (incl. 95% CI)  | Both Sexes             | Males                  | Females                |
|---|------------------------|------------------------|------------------------|
| <b>Step 1 Cervical Cancer Screening</b>   |                        |                        |                        |
| Percentage of women aged 30-49 years who have ever had a screening test for cervical cancer   |                        |                        | 15.5<br>(13.4-17.6)    |
| <b>Step 2 Physical Measurements</b>   |                        |                        |                        |
| Mean body mass index - BMI (kg/m <sup>2</sup> )   | 26.4<br>(26.2-26.5)    | 25.3<br>(25.1-25.5)    | 27.5<br>(27.5-27.8)    |
| Percentage who are overweight (BMI ≥ 25 kg/m <sup>2</sup> )   | 55.6<br>(53.9-57.2)    | 48.3<br>(46.1-50.6)    | 63.3<br>(61.4-65.2)    |
| Percentage who are obese (BMI ≥ 30 kg/m <sup>2</sup> )  | 21.8<br>(20.5-23.1)    | 14.1<br>(12.7-15.5)    | 30.1<br>(28.3-31.9)    |
| Average waist circumference (cm)  |                        | 92.2<br>(91.3-93.1)    | 92.8<br>(91.9-93.7)    |
| Mean systolic blood pressure - SBP (mmHg), including those currently on medication for raised BP  | 126.3<br>(125.6-127.0) | 127.7<br>(126.9-128.5) | 125.0<br>(124.1-125.8) |
| Mean diastolic blood pressure - DBP (mmHg), including those currently on medication for raised BP   | 75.1<br>(74.7-75.6)    | 74.9<br>(74.3-75.4)    | 75.4<br>(74.8-75.9)    |
| Percentage with raised BP (SBP ≥ 140 and/or DBP ≥ 90 mmHg or currently on medication for raised BP)   | 23.6<br>(22.2-25.0)    | 23.1<br>(21.3-24.9)    | 24.1<br>(22.4-25.7)    |
| Percentage with raised BP (SBP ≥ 140 and/or DBP ≥ 90 mmHg) who are not currently on medication for raised BP  | 71.9<br>(69.4-74.4)    | 80.8<br>(77.8-83.8)    | 63.2<br>(59.8-66.6)    |
| <b>Step 3 Biochemical Measurement</b>   |                        |                        |                        |
| Mean fasting blood glucose, including those currently on medication for raised blood glucose [mg/dl]  | 98.0<br>(96.7-99.3)    | 97.6<br>(96.0-99.3)    | 98.4<br>(96.7-100.1)   |
| Percentage with impaired fasting glycaemia as defined below<br>• plasma venous value ≥ 6.1 mmol/L (110 mg/dl) and < 7.0 mmol/L (126 mg/dl)<br>• capillary whole blood value ≥ 5.6 mmol/L (100 mg/dl) and < 6.1 mmol/L (110 mg/dl)   | 8.2<br>(7.1-9.3)       | 8.1<br>(6.7-9.4)       | 8.3<br>(7.0-9.6)       |
| Percentage with raised fasting blood glucose as defined below or currently on medication for raised blood glucose<br>• plasma venous value ≥ 7.0 mmol/L (126 mg/dl)<br>• capillary whole blood value ≥ 6.1 mmol/L (110 mg/dl)   | 9.0<br>(8.1-9.9)       | 8.4<br>(7.2-9.7)       | 9.6<br>(8.5-10.7)      |
| Percentage Blood glucose ≥ 126. or HbA1c ≥ 6.5. or takes meds   | 14.4<br>(12.6-16.3)    | 14.4<br>(12.0-16.0)    | 14.9<br>(12.9-17.0)    |
| Mean total blood cholesterol, including those currently on medication for raised cholesterol [mg/dl]  | 162.9<br>(161.3-164.4) | 159.2<br>(157.4-161.0) | 166.6<br>(164.7-168.5) |
| Percentage with raised total cholesterol (≥ 5.0 mmol/L or ≥ 190 mg/dl or currently on medication for raised cholesterol)  | 24.0<br>(22.5-25.5)    | 21.5<br>(19.7-23.2)    | 26.6<br>(24.6-28.5)    |
| Mean intake of salt per day (in grams)  | NA                     | NA                     | NA                     |
| <b>Cardiovascular disease (CVD) risk</b>  |                        |                        |                        |
| Percentage aged 40-69 years with a 10-year CVD risk ≥ 30%, or with existing CVD**   | 8.7<br>(7.4-10.0)      | 8.9<br>(7.3-10.6)      | 8.5<br>(6.9-10.2)      |
| <b>Summary of combined risk factors</b>   |                        |                        |                        |
| <ul style="list-style-type: none"> <li>current daily smokers</li> <li>less than 5 servings of fruits &amp; vegetables per day</li> <li>insufficient physical activity</li> <li>overweight (BMI ≥ 25 kg/m<sup>2</sup>)</li> <li>raised BP (SBP ≥ 140 and/or DBP ≥ 90 mmHg or currently on medication for raised BP)</li> </ul> |                        |                        |                        |
| Percentage with none of the above risk factors  | 3.4<br>(2.6-4.2)       | 3.6<br>(2.6-4.7)       | 3.2<br>(2.3-4.1)       |
| Percentage with three or more of the above risk factors, aged 18 to 44 years  | 22.7<br>(21.1-24.3)    | 22.0<br>(19.9-24.1)    | 23.4<br>(21.0-25.8)    |
| Percentage with three or more of the above risk factors, aged 45 to 69 years  | 45.0<br>(42.4-47.6)    | 39.1<br>(36.0-42.1)    | 51.0<br>(47.7-54.3)    |
| Percentage with three or more of the above risk factors, aged 18 to 69 years  | 29.6<br>(27.9-31.2)    | 27.2<br>(25.2-29.1)    | 32.1<br>(29.9-34.4)    |

\*\* A 10-year CVD risk of ≥ 30% is defined according to age, sex, blood pressure, smoking status (current smokers OR those who quit smoking less than 1 year before the assessment), total cholesterol, and diabetes (previously diagnosed OR a fasting plasma glucose concentration > 7.0 mmol/l (126 mg/dl)).

**please contact: STEPS country focal point : Dr.Nadir-Azirou Djamilia**  
**Azirounadir@yahoo.fr**