

SKY Sokhapeap Kon Youeng - Our Child Health - Prevention or treatment of moderate malnutrition - Infants and young children|Lactating women (LW)|Pregnant women (PW)

Programme Data

Action data

Start date	
Country(ies):	Cambodia
Status:	On-going
Area:	Rural
Place:	Banteay Meanchey, 3 health centers, 6 villages
Topic:	Management of moderate malnutrition
Target group:	Infants and young children Lactating women (LW) Pregnant women (PW)
Age group:	0-59 months
Delivery:	Community-based Primary health care center
Implementation details :	<p>5 staff</p>
Target population size :	128,234
Coverage level (%):	23%
Outcome indicator(s):	<ul style="list-style-type: none"> % of children 0-23 months who were put to the breast within one hour of birth % of infants (0-5 months of age) exclusively breast-fed during past 24 hours % of children 6-23 months of age who received minimum meal frequency during the previous day % of children 6-23 months of age who were fed with minimum dietary diversity (four or more food groups) during the previous day Proportion of children 0-23 months with diarrhea in the past two weeks, who were treated with oral rehydration solution (ORS), zinc and appropriate feeding % of mothers of children age 0-23 months who had four or more antenatal visits during last pregnancy (as verified by ANC record) % of mothers with children 0-23 months of age consumed at least 90 iron/folic acid tablets during last pregnancy % of mothers with children 0-23 months who delivered their baby with a skilled birth attendant % of mothers with children 0-23 months of age consumed at least 42 iron/folic acid tablets during the first six weeks after the delivery of the youngest child % of mothers with children 0-23 months of age who received at least two post-natal visits from a trained health care worker during the first week after the birth of their youngest child % of children age 12-23 months who have completed 3rd DTP dose plus measles vaccination
M&E system:	<ul style="list-style-type: none"> Household survey (with anthropometry 0-23 months) 0-59 months LQAS Register Child 0-36 months Weighing every 2 months Home visit
Baseline:	Stunting under fives 23.2%Wasting under fives 8.9%Underweight under fives 18.3%
Post-intervention:	N/A
Outcome reported by social determinants:	Vulnerable groups
Typical problems	Solutions
Insufficient staff	