

Declaration of Labelling

1. Product Identification

Product name: KLYPSON 500 WG

This product is Water Dispersible Granules (WG) packed in 150g metalized aluminium pouch.

Manufactured and distributed by:

Tagros Chemicals India Private Limited
"Jhaver Centre", Rajah Annamalai Building,
IV Floor, 72, Marshalls Road, Egmore,
Chennai-600 008, India.

Phone no: +91-44-42007400.

Email: info@tagros.com

Website: www.tagros.com

2. Ingredient statement

Clothianidin (#210880-92-5): 50%

Other ingredients: 50%

Clothianidin is a neonicotinoids in IRAC group 4A.

3. Safety

Precaution:

IF ON SKIN: Wash with plenty of water and soap

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

IF INHALED: Remove person to fresh air and keep comfortable for breathing Immediately call a POISON CENTER or doctor Wash hands thoroughly after handling Wear protective gloves .Avoid release to the environment

Warning:

Keep out of reach of children.

Harmful if swallowed

May Causes Allergic skin irritation

Very toxic to aquatic life with long lasting effects.

First aid:

Eye Contact: Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Get medical attention if irritation develops and persists.

Skin Contact: Wash off immediately with soap and plenty of water while removing all contaminated clothes and shoes. Get medical attention if irritation develops and persists.

Inhalation: Move person from contaminated area to fresh air. Seek medical attention immediately. If person is not breathing, give artificial respiration, preferably by mouth-to-mouth, if possible.

Ingestion: Do not induce vomiting without medical advice. Rinse mouth with water and afterwards drink plenty of water or milk. If vomiting does occur, rinse mouth and drink fluids again. Never give anything by mouth to an unconscious person. Immediate medical attention is required.

4. Direction for use

This product is only intended to be used as an Indoor Residual Spray. It has been shown to control Anopheline mosquitoes which are known to transmit malaria.

Apply to indoor walls, eaves, and other surfaces on which mosquitoes or other target vectors may rest.

Do not apply to surfaces used for food preparation or eating.

Target application rate:

The target dose of the product is 300 mg a.i./m². In order to achieve the target dose, it is important to follow the labelled directions or dilution, mixing and spraying instructions.

Equipment:

This product should only be applied with hand-held compression sprayers complying with WHO Specifications, fitted with flat fan nozzles under low to medium pressure. Constant Flow Valves can be used. Ensure that the spraying equipment functions correctly, is well maintained and does not leak.

Spray preparation/Mixing instructions:

Fill the tank half full of clean or filtered water. For 150g sachets, add the contents of one sachet to the tank. Fill the tank with clean or filtered water to the appropriate volume. Seal the tank, pressurize and mix the contents of the tank by inverting several times.

Prepare only the amount of required diluted product for immediate usage. Do not store diluted material.

Dilution table:

Nozzle type	Sachet	Required volume of water (L)	Surface area (m ²) to be covered
8002E (with Constant Flow Valve)	150g	7.5	250
8002E (without Constant Flow Valve)	150g	10	250

Instructions for Spraying

Pressurize the tank in accordance with the manufactures instructions. The tank should be pressurized to 55 psi when using 8002E nozzle type. The target pressure should be maintained throughout the application. If pressure drops below 35 psi, re-pressurize the tank and continue spraying.

Apply spray in vertical swathes 75 cm wide, with an overlap of 5 cm. Spray from roof to floor, using a downward motion, to complete one swathe. Step sideways and spray upwards from floor to roof.

To ensure the correct swathe width, keep the spray tip about 45 cm from the wall. Lean forwards as you spray from top of the wall and move back as you bring the nozzle downwards, spraying at a tempo of 2.5 seconds per metre of surface sprayed.

Post-Spray application

Do not allow people or pets to enter the treated dwelling until sprays have dried. Floors should be cleaned before allowing children and pets to enter the dwelling.

5. Product claims

To control Anopheline mosquitoes which are known to transmit Malaria.