

COMIRNATY® (Omicron JN.1)

COVID-19 mRNA Vaccine (nucleoside modified)

Formulation	Grey Cap	Orange Cap	Blue Cap	Maroon Cap	Yellow Cap
Dosage	30 µg	10 µg	10 µg	3 µg	3 µg
Presentation	Single and Multi-dose vial	Multi-dose vial	Single and Multi-dose vial	Multi-dose vial	Multi-dose vial
Pharmaceutical Form	Dispersion for injection	Concentrate for dispersion for injection	Dispersion for injection	Concentrate for dispersion for injection	Concentrate for dispersion for injection
Number of doses	1 and 6	10	1 and 10	10	3
Dilution	Dilution NOT required	Dilution required	Dilution NOT required	Dilution required	Dilution required
Diluent	NOT required	Sodium Chloride Inj. USP 0.9%	NOT required	Sodium Chloride Inj. USP 0.9%	Sodium Chloride Inj. USP 0.9%
Route of Administration	Intramuscular	Intramuscular	Intramuscular	Intramuscular	Intramuscular
Age Indication	12 years and older	Children from 5 to 11 years	Children from 5 to 11 years	Children between 6 months to 4 years	Children between 6 months to 4 years
Vaccine Vial Monitor	None	None	None	None	None
Vaccine Preservative	None	None	None	None	None
Refrigeration Storage Time (2 °C to 8 °C)	10 weeks	10 weeks	10 weeks	10 weeks	10 weeks
Shelf-life at ULT Freezer Storage Time (-90 °C to -60 °C)	18 months	18 months	18 months	18 months	18 months
Multidose Vial Policy	WHO recommends that opened vials of this vaccine should be discarded 6 hours after opening or at the end of the immunization session, whichever comes first.				

منظمة الصحة العالمية • 世界卫生组织

Organisation mondiale de la Santé • Всемирная организация здравоохранения • Organización Mundial de la Salud