

COMIRNATY® (Omicron JN.1)

COVID-19 mRNA Vaccine (nucleoside modified)

Formulation	Grey Cap	Grey Cap new	Orange Cap	Blue Cap	Maroon Cap	Yellow Cap
Dosage	30 µg	30 µg	10 µg	10 µg	3 µg	3 µg
Presentation	Single and Multi-dose vial	Multi-dose vial	Multi-dose vial	Single and Multi-dose vial	Multi-dose vial	Multi-dose vial
Pharmaceutical Form	Dispersion for injection	Dispersion for injection	Concentrate for dispersion for injection	Dispersion for injection	Concentrate for dispersion for injection	Concentrate for dispersion for injection
Number of doses	1 and 6	6	10	1 and 10	10	3
Dilution	Dilution NOT required	Dilution NOT required	Dilution required	Dilution NOT required	Dilution required	Dilution required
Diluent	NOT required	NOT required	Sodium Chloride Inj. USP 0.9%	NOT required	Sodium Chloride Inj. USP 0.9%	Sodium Chloride Inj. USP 0.9%
Route of Administration	Intramuscular	Intramuscular	Intramuscular	Intramuscular	Intramuscular	Intramuscular
Age Indication	12 years and older	12 years and older	Children from 5 to 11 years	Children from 5 to 11 years	Children between 6 months to 4 years	Children between 6 months to 4 years
Vaccine Vial Monitor	None	None	None	None	None	None
Vaccine Preservative	None	None	None	None	None	None
Refrigeration Storage Time (2 °C to 8 °C)	10 weeks	12 months (shelf life)	10 weeks	10 weeks	10 weeks	10 weeks
Shelf-life at ULT Freezer Storage Time (-90 °C to -60 °C)	18 months	DO NOT FREEZE	18 months	18 months	18 months	18 months
Multidose Vial Policy	WHO recommends that opened vials of this vaccine should be discarded 6 hours after opening or at the end of the immunization session, whichever comes first.					

منظمة الصحة العالمية • 世界卫生组织

Organisation mondiale de la Santé • Всемирная организация здравоохранения • Organización Mundial de la Salud