

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

Artesun 30mg*

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each vial contains: artesunate powder 30 mg

Each ampoule of solvent contains: sodium bicarbonate 50 mg/ml, 0.5 ml

Each ampoule of diluent contains: sodium chloride 9 mg/ml, 2.5 ml

3. PHARMACEUTICAL FORM

Artesunate for injection: White crystalline powder

Solvent (sodium bicarbonate injection): Clear, colourless liquid

Diluent (sodium chloride injection): Clear, colourless liquid

4. CLINICAL PARTICULARS

4.1 Therapeutic indication

Artesun® 30mg, administered intravenously or intramuscularly, is indicated for the treatment of severe malaria caused by *Plasmodium falciparum*, in adults and children.

Consideration should be given to official treatment guidelines for malaria (e.g. by WHO).

4.2 Posology and method of administration

Dose:

Adults and children weighing more 20 kg or more: **Artesun® 30mg** is administered at a dose of 2.4 mg of artesunate / kg body weight, by intravenous (IV) or intramuscular (IM) injection, at 0, 12 and 24 hours, then once daily until oral treatment can be substituted.

Children weighing less than 20 kg: **Artesun® 30mg** is administered at a dose of 3 mg of artesunate / kg body weight, by intravenous (IV) or intramuscular (IM) injection, at 0, 12 and 24 hours, then once daily until oral treatment can be substituted (see section 5.1).

Method of administration

Artesun® 30mg should be administered for a minimum of 24 hours (3 doses), regardless of the patient's ability to tolerate oral medication earlier. After at least 24 hours of **Artesun® 30mg**, and when able to tolerate oral medication, the patient should be switched to a complete treatment course of an oral combination antimalarial regimen. Relevant treatment guidelines should be consulted when selecting an appropriate regimen, e.g. those of the WHO (see reference section at end of this SmPC).

* Trade names are not prequalified by WHO. This is the national medicines regulatory authority's responsibility. Throughout this WHOPAR the proprietary name is given as an example only.

Preparation

Because of the instability of artesunate in aqueous solutions the reconstituted solution must be used within one hour of preparation. Therefore the required dose of artesunate should be calculated (dose in mg = patient's weight in kg x 2.4 or dose in mg = patient's weight in kg x 3 for children weighing less than 20 kg, respectively) and the number of vials of artesunate needed should be determined prior to reconstituting the artesunate powder.

Reconstitution of the artesunate solution

Using a syringe, withdraw 0.5 ml of the supplied sodium bicarbonate solvent from the ampoule and inject into the vial containing the artesunate powder. Shake the vial for several minutes to mix well until the powder is completely dissolved and the solution is clear. If the solution appears cloudy or a precipitate is present, it should be discarded. The reconstituted artesunate solution should always be used immediately and discarded if not used within one hour.

Following reconstitution the solution must be diluted according to the method of injection, as described below.

For intravenous (IV) injection (10 mg/ml)

Using a syringe, add 2.5 ml of sodium chloride 0.9% for injection to the vial containing the reconstituted artesunate solution. This will yield 3 ml of a solution containing artesunate 10 mg/ml. Shake to mix well, ensuring that the resulting solution is still clear. If the solution appears cloudy or a precipitate is present, it should be discarded.

The volume of the solution required (ml) will be:

$$\text{Volume (ml)} = [\text{dose (mg)}] \div 10$$

Withdraw the required volume of artesunate solution from the vial with a syringe and then inject slowly intravenously, over 1-2 minutes.

Artesun® 30mg should NOT be administered as an intravenous drip.

For intramuscular (IM) injection (20 mg/ml)

Using a syringe, add 1 ml of sodium chloride 0.9% for injection to the vial containing the reconstituted artesunate solution. This will yield 1.5 ml of a solution containing artesunate 20 mg/ml. Shake to mix well, ensuring that the resulting solution is still clear. If the solution appears cloudy or a precipitate is present, it should be discarded.

The volume of the solution required (ml) will be:

$$\text{Volume (ml)} = [\text{dose (mg)}] \div 20$$

Withdraw the required volume of artesunate solution from the vial with a syringe and then inject intramuscularly; the anterior thigh is usually the preferred site for injection. If the total volume of solution to be injected intramuscularly is large, it may be preferable to divide the volume and inject it at several sites, e.g. both thighs.

Do not use water for injection for reconstitution of the artesunate powder or for dilution of the resulting solution prior to injection.

Hepatic and renal impairment:

Dose adjustment is not necessary in patients with hepatic or renal impairment (see Sections 4.4 and 5.2).

4.3 Contraindications

Artesun® 30mg is contraindicated in patients with hypersensitivity to artesunate or other artemisinins.

4.4 Special warnings and precautions for use

Non-falciparum malaria

Artesunate has not been evaluated in the treatment of severe malaria due to *Plasmodium vivax*, *Plasmodium malariae* or *Plasmodium ovale*.

Resistance to antimalarials

Local information on the prevalence of resistance to antimalarials should be considered in choosing the appropriate combination antimalarial regimen for use with **Artesun® 30mg**. Relevant treatment guidelines should be consulted such as those of the WHO and public health authorities (see reference section at end of this SmPC)

Post-treatment haemolytic anaemia

Delayed haemolytic anaemia following treatment with injectable artesunate has been observed in children and adults in malaria endemic areas and in non-immune travelers presenting with severe falciparum malaria. The risk was most pronounced in patients with hyperparasitaemia and in younger children. Some cases have been severe and required blood transfusion. Vigilance for delayed onset anaemia is therefore advised, particularly in hyperparasitaemic patients and younger children. Given the persistent uncertainties regarding the risk of delayed hemolysis, a minimum follow-up should be strictly observed until day 28. (See Section 4.8 and References)

Hepatic / renal impairment:

Data regarding artesunate pharmacokinetics in patients with hepatic and/or renal impairment are limited. Based on data from studies in patients with severe malaria, as well as the known metabolism of artesunate (see Section 5.2), dosage adjustment is not considered necessary in patients with hepatic or renal impairment.

Paediatric population

In clinical trials, the efficacy and safety of intravenous and intramuscular artesunate have been similar in adult and paediatric populations.

4.5 Interaction with other medicinal products and other forms of interaction

Artesunate is rapidly and extensively converted to dihydroartemisinin (DHA), the active metabolite, primarily by plasma and erythrocyte esterases. DHA elimination is also rapid (half-life approximately 45 min) and the potential for drug-drug interactions appears limited. *In vitro* drug-interaction studies

have demonstrated minimal effects of artesunate on cytochrome P450 isoenzymes. Few clinical drug-drug interaction studies have been performed, however no clinically significant interactions have been identified.

4.6 Pregnancy and lactation

Pregnancy

Severe malaria is especially hazardous during pregnancy, therefore full dose parenteral artesunate treatment should be administered at any stage of pregnancy without delay.

In animal studies, artesunate has been associated with fetal toxicity during the first trimester of pregnancy. Limited clinical experience with the use of artesunate in the first trimester of pregnancy as well as clinical data from pregnant women treated with artemisinin derivatives in the second and third trimester, do not indicate adverse effects of artesunate on pregnancy or on the health of the fetus /newborn child. In addition, an observational study conducted in 1179 pregnant women with first-trimester falciparum malaria (of which 183 were treated with first line artemisinin derivatives) showed no differences in the risk of miscarriage or major congenital malformation in artemisinin derivatives versus quinine treatment. (Moore KA 2016, see References.)

Breast-feeding

Limited information indicates that dihydroartemisinin, the active metabolite of artesunate, is present at low levels in breast milk. The drug levels are not expected to cause any adverse effects in breastfed infants. The amount of drug present in breast milk does not protect the infant from malaria.

Fertility

No specific studies with artesunate in humans have been conducted to evaluate effects on fertility. In a reproduction toxicity study in rats, testicular and epididymal lesions were seen, but there were no effects on fertility (see section 5.3). The relevance of this finding for humans is unknown.

4.7 Effects on ability to drive and use of machines

There is no information on the effect of artesunate on the ability to drive or use machines. The patient's clinical status should be considered when assessing ability to drive or operate machinery.

4.8 Undesirable effects

The most important reported side effect of artesunate is a rare severe allergic reaction (estimated risk approximately 1 in 3000 patients), which has involved urticarial rash as well as other symptoms, including hypotension, pruritus, oedema, and/or dyspnoea. A potentially serious delayed hemolysis (Post-Artesunate Delayed Hemolysis, PADH) has been reported frequently in travelers and in children. See below. *

More common minor side effects associated with IV administration have included dizziness, light-headedness, rash, and taste alteration (metallic/ bitter taste). Transient elevation in liver transaminases, nausea, vomiting, anorexia and diarrhea have also been commonly reported, however it is uncertain whether such events may have been symptoms of severe malaria.

Adverse events considered at least possibly related to artesunate are listed below by body system, organ class and absolute frequency. Frequencies are defined as very common ($\geq 1/10$), common

(1/100–1/10), uncommon (1/1000–1/100), rare (1/10 000–1/1000), and very rare (< 1/10 000).

Blood and lymphatic systems disorders

Very common: Post-treatment haemolytic anaemia in travelers, mild and transient decrease in reticulocyte count (See References)

Common to very common: Post-treatment haemolytic anaemia in endemic areas

Uncommon: Neutropenia and anaemia (both occasionally severe), thrombocytopenia, agranulocytosis, reticulocytopenia, erythroblastopenia

Very rare: Pure red cell aplasia

Nervous system disorders

Common: Dizziness, light-headedness, headache, insomnia, tinnitus (with or without decrease in auditory function)

Very rare: Peripheral neuropathy (or paraesthesia)

Cardiovascular disorders

Uncommon: Rhythm and conduction disorders. Some cases of cardiac adverse events and particularly rhythm (bradycardia, sinus arrhythmia) and conduction disorders (QTc lengthening, abnormal sinoatrial conduction) have been recently described. One case of cardiac arrest has been reported in a context of severe malaria with multi-organ failure but the causality by Artesunate has not been established. Whether or not cardio-vascular events should be attributed to artesunate or to severe malaria is not known, but considering that other artemisinin derivatives (arthemeter, arteether) have been associated with QT prolongation in pre-clinical data, EKG should be monitored before and during treatment, especially in patients with a history of cardiovascular impairment or having risk factors.

Rare: Arterial ischemia, hypertensive retinopathy

Respiratory disorders

Common: Cough and/ or nasal symptoms

Gastrointestinal disorders

Common: Altered taste, nausea, vomiting, abdominal pain or cramps, diarrhoea

Rare: Raised serum amylase, pancreatitis

Hepatobiliary disorders

Uncommon: Transient rises in liver transaminases (AST, ALT)

Rare: Hepatitis, calculous cholecystitis

Skin and subcutaneous tissue disorders

Common: Rash, alopecia

Musculoskeletal and connective tissue disorders

Common: Arthralgia, muscle disorders

General disorders and administration site conditions

Common: Fatigue, malaise, fever, pain at injection site

Immune system disorders

Uncommon: hypersensitivity

**Post-Artesunate Delayed Hemolysis (PADH)*

A delayed hemolysis was detected in prospective studies with a frequency ranging from 7 to 27% of patients treated with artesunate. This adverse event has been observed both in travellers (15%) and among young African children (7-22%) in cases of severe malaria (See References section). The best known cases of delayed haemolysis have been reported in travellers (including non-immune tourists and “visiting friends and relatives” persons). The pathophysiology of this phenomenon has not been fully elucidated, but may include various combinations of delayed destruction of different subpopulations of erythrocytes and an immune-mediated etiology. All patients treated with parenteral artesunate should be followed for at least 4 weeks to detect signs of hemolysis and enable appropriate treatment.

Background

Definition: The current definition of PADH is the association of a drop in hemoglobin levels of more than 10% and a rise in lactate dehydrogenase levels greater than 10 %, both occurring more than 8 days after starting treatment with artesunate.

Incidence: Reported incidence ranges between 0% and 63 %. A recent systematic analysis (Roussel et al, 2017) found 15.3% (74/485) of patients to be affected by delayed hemolysis. Exactly half of PADH patients with available data were transfused (27/54 patients). Assessment in the single prospective study available so far (Jaureguiberry et al, 2015) reported that 27% (21/78) of patients with follow-up data beyond Day 8 had PADH per the current definition. The nadir (lowest value) in hemoglobin during follow-up was lower than 7 g/dL in 15% of patients with PADH. One of those patients required a transfusion. No sequelae or death have been reported in patients with PADH so far, but in one report in an endemic area (Plewes et al, 2015) a patient had acute renal insufficiency probably related to the acute hemolytic event with hemoglobinuria, and one African child had coma probably related to PADH (Rolling et al, 2014). Delayed hemolytic episodes have not been reported in Phase 3 clinical trials in endemic areas possibly due to the lack of precise follow-up between Day 3-7 and Day 28 in these trials. An incidence of PADH of 7% was observed in a prospective study of 72 African children treated with artesunate for severe malaria. Parameters associated with PADH were younger age and higher initial parasitemia. PADH affected 11.4% of patients (children and adults) in the Democratic Republic of Congo. All episodes resolved rapidly. The impact of the declining incidence of malaria and waning antimalarial immunity in endemic countries on the incidence and impact of PADH is not known, although these parameters have an influence on the mechanism of parasite clearance, related itself to the mechanism of PADH.

Impact: PADH and other post-artesunate hemolytic episodes almost invariably resolve in less than a few weeks, often in a few days. Transfusion indication is per current standard of care. Whether long term sequelae occur in a small proportion of patients is not known. The overall risk/benefit remains highly favourable for injectable artesunate, with a large mortality advantage over quinine. WHO strongly recommends continued use of artesunate for severe malaria.

Paediatric population:

The safety profile of injectable artesunate is similar in children and adults.

4.9 Overdose

Experience of acute overdose with artesunate is limited. A case of overdose has been documented in a

5-year-old child who was inadvertently administered rectal artesunate at a dose of 88 mg/kg/day over 4 days, representing a dose more than 7-fold higher than the highest recommended artesunate dose. The overdose was associated with pancytopenia, melena, seizures, multi-organ failure and death.

Treatment of overdose should consist of general supportive measures.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antimalarial , ATC code: P01BE03

Mechanism of action

Artesunate is a hemisuccinate derivative of dihydroartemisinin, which is itself formed by the reduction of artemisinin. Artemisinin is a sesquiterpene lactone endoperoxide extracted from qinghao (sweet wormwood, *Artemisia annua L.*), a plant which has been used for centuries in traditional Chinese medicine.

The mechanism of action of the artemisinins likely involves cleavage of the internal endoperoxide bridge through reaction with haeme within the infected erythrocyte, thereby generating free radicals which alkylate vital parasite proteins. However, artemisinins have also been reported to inhibit an essential parasite calcium adenosine triphosphatase.

The artemisinins are distinguished from other antimalarials by their ability to kill all erythrocytic stages of the malaria parasite, including the relatively inactive ring stage and late schizonts, as well as the gametocytes responsible for malaria transmission. Artesunate and the artemisinins are the most rapid acting of the antimalarials, and they have also been shown to enhance splenic clearance of infected erythrocytes by reducing cytoadherence.

In vitro, dihydroartemisinin (DHA), the active metabolite of artesunate, exhibits similar potency against chloroquine-resistant and chloroquine-sensitive clones of *P. falciparum*.

Artesunate and the other artemisinins are essentially inactive against extra-erythrocytic forms, sporozoites, liver schizonts or merozoites.

Clinical efficacy and safety

In the SEAQUAMAT (South East Asian Quinine Artesunate Malaria Trial), an international randomised, open-label, multicenter trial conducted in Bangladesh, India, Indonesia and Myanmar, 1461 patients with severe malaria (including 1259 adults) were treated intravenously with either artesunate or quinine. Artesunate was administered at 2.4 mg/kg IV at 0, 12 and 24 h and then every 24 h until the patient could tolerate oral medication. Quinine was given IV at 20 mg/kg over 4 hours, followed by 10 mg/kg over 2-8 hours, 3 times daily until oral therapy could be started. Mortality in the artesunate group was 15% versus 22% in the quinine group, for a reduction in risk of death of 34.7% (p=0.0002). Subgroup analysis suggested a greater benefit of artesunate versus quinine in

patients with parasitaemia >10%. The reduction in mortality observed in the 202 paediatric patients (<15 years of age) appeared consistent with the overall results, however the number of children was too small to demonstrate statistical significance. Post-treatment hypoglycaemia was more common in the quinine-treated group.

Paediatrics

The AQUAMAT (African Quinine Artesunate Malaria Trial) was an international, randomized open-label multicenter trial which sought to extend the results of the SEAQUAMAT study by comparing parenteral artesunate versus IV quinine for severe malaria in 5425 African children (< 15 years) in 9 African countries (Mozambique, The Gambia, Ghana, Kenya, Tanzania, Nigeria, Uganda, Rwanda, and Democratic Republic of the Congo). Dosing was similar to SEAQUAMAT, except that both artesunate and quinine could be administered either intravenously or intramuscularly, using the same doses for IM and IV administration for each drug. Roughly one third of patients received study drug by intramuscular injection. Mortality in the artesunate group was 8.5% compared to 10.9% in the quinine group, resulting in a relative risk reduction for death of 22.5% ($p=0.0022$); the risk reduction was similar for IV and IM administration. In addition, although the risk of neurological sequelae in survivors in both groups did not differ significantly by 28 days following treatment, in-hospital coma, convulsions, and deterioration of coma were all less frequent in the artesunate-treated patients. As in the SEAQUAMAT, post-treatment hypoglycaemia was more common in the quinine-treated group.

5.2 Pharmacokinetic properties

Intravenous

After intravenous injection artesunate is very rapidly biotransformed to its active metabolite, dihydroartemisinin (DHA). Consequently, artesunate half-life ($t_{1/2}$) is estimated to be less than 5 minutes. Following a single IV dose of 2.4 mg/kg, maximum artesunate plasma concentrations (C_{max}) were estimated to be 77 $\mu\text{mol/L}$ in a study in Gabonese children with severe malaria, and 42 and 36 $\mu\text{mol/L}$ in two studies in Vietnamese adults with uncomplicated malaria.

High concentrations of DHA are observed within 5 minutes of artesunate IV administration. In the above studies (adult and paediatric), the ranges of values for the estimated time to maximum concentration (t_{max}) and $t_{1/2}$ for DHA were 0.5-15 minutes and 21-64 minutes, respectively, while DHA C_{max} values ranged from 5.3-10.6 $\mu\text{mol/L}$.

Intramuscular

Artesunate is rapidly absorbed following intramuscular injection, and peak plasma levels are generally achieved within 30 minutes of administration. Thus, after IM injection of 2.4 mg/kg of artesunate, absorption was rapid in Gabonese children and Vietnamese adults, with T_{max} values of 8 and 12 minutes, respectively. The corresponding artesunate $t_{1/2}$ values were estimated to be 48 minutes in children and 41 minutes in adults, and C_{max} values were 1.7 and 2.3 $\mu\text{mol/L}$, for children and adults, respectively.

After IM injection artesunate C_{max} values were therefore lower by roughly 45-fold in children and 20-fold in adults when compared to IV injection. However, rates of artesunate elimination in children and adults were 32-fold and 13-fold slower, respectively, following IM injection, compared to IV administration.

Distribution

DHA has been shown to substantially accumulate in *P. falciparum*-infected erythrocytes. Plasma protein binding of dihydroartemisinin was determined to be 93% in patients and 88% in healthy volunteers.

Metabolism and elimination

Artesunate is extensively and rapidly hydrolysed by plasma esterases, with possible minimal contribution by CYP2A6. The main metabolite, dihydroartemisinin, accounts for most of the *in vivo* antimalarial activity of oral artesunate, however, following IV administration, artesunate may contribute more significantly. DHA is further metabolized in the liver via glucuronidation and is excreted in the urine; α -dihydroartemisinin- β -glucuronide has been identified as the major urinary product in patients with *falciparum* malaria.

Special population:

No pharmacokinetic data are available for patients with impaired renal or hepatic function. However, based on the known mechanisms of metabolism and elimination of artesunate, combined with clinical data from patients with severe malaria and accompanying renal and/or hepatic compromise of various degrees, no dose modifications are considered necessary in renal or hepatic impairment.

5.3 Preclinical safety data

General toxicity

Artesunate presents low acute toxicity. After repeated administration of 50 mg/kg/day in rats and 82.5 mg/kg/day in dogs, i.e. approximately 10 and 17 times the proposed maximal therapeutic dose in man, rare occurrences of moderate hepatotoxicity (4%) were reported, with regression in 100% of cases without sequelae. Hematopoietic toxicity on the red and white line is responsible for reticulocytopenia and leucopenia of central origin. The toxicity appears to be on the progenitors of the erythroid lineage. There is also a negative effect on the phagocytosis capacities of neutrophils and inhibition of lymphocyte proliferation (in vitro).

Genotoxicity

Artesunate did not show any mutagenic or clastogenic potential in *in vitro* and *in vivo* tests (Ames, mouse micronucleus).

Carcinogenesis

No studies of the carcinogenic potential of artesunate have been conducted.

Reproductive toxicology studies

Oral artesunate caused dose-dependent foetal toxicity in rats, rabbits and monkeys, resulting in foetal resorption and abortion, as well as a low incidence of cardiac and skeletal defects. The no-observed-adverse-effect-level (NOAEL) was 12 mg/kg in pregnant monkeys (3 and 7 day exposures) and the no or low adverse effects level was 5-7 mg/kg in pregnant rats or rabbits (12 day exposures), both of which are above the therapeutic dose range (2.4-4.8 mg/kg) and expected duration

of exposure for treatment of severe malaria in humans. In rats, the embryo-fetuses were most sensitive from gestational days 9-14; at other times embryotoxicity was significantly reduced. A study of artesunate administered to male rats daily for 6 weeks noted testicular and epididymal lesions, although these lesions did not affect fertility. The lesions were reversible after cessation of treatment.

Safety pharmacology studies

A slight sedative effect, decrease in body temperature, mild natriuretic effect and a decrease in creatinine clearance were observed with artesunate after single intravenous doses of 200 mg/kg (mice), 450 mg/kg (rats, rabbits and dogs) and following single oral doses of 180 mg/kg in male rats. Beagle dogs administered IV artesunate at 10, 20, 50, and 50 mg/kg for 14 days did not display significant clinical effects, including any signs of neurotoxicity, effects on body weight, ECG abnormalities (including QT interval changes), heart rate, blood pressure, or respiratory rate.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Artesunate powder for injection: no excipients

Solvent: sodium bicarbonate and water for injection

Diluent: sodium chloride and water for injection

6.2 Incompatibilities

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

6.3 Shelf life

36 months

6.4 Special precautions for storage

Store below 30°C. Protect from light.

The reconstituted solution should be stored below 30°C and should be used within 1 hour.

6.5 Nature and contents of container

Artesunate for injection: The primary packs are colourless 7 ml, type I glass vials with gray colored type I rubber stoppers, crimped with aluminum-plastic cap.

Solvent (sodium bicarbonate injection 50mg/ml): The primary packs are colourless type I glass ampoules of 1ml.

Diluent (sodium chloride injection 9mg/ml): The primary packs are colourless type I glass ampoules of 5ml.

Pack size: A small box containing one vial of artesunate for injection, one ampoule of the sodium bicarbonate injection (solvent) and one ampoule of the sodium chloride injection (diluent).

6.6 Special precautions for disposal

No special requirements.

7. SUPPLIER

Guilin Pharmaceutical Co., Ltd.;
No. 43 Qilidian Road,
Guilin,
Guangxi, China
Telephone: + 86 773 3675053
Fax: +86 773 2670530
Email: glpharma@public.glptt.gx.cn

8. WHO REFERENCE NUMBER (PREQUALIFICATION PROGRAMME)

MA089

9. DATE OF FIRST PREQUALIFICATION

May 2013

10. DATE OF REVISION OF THE TEXT

November 2015
Section 4 and 5 were updated in January 2020.

Reference list:

General

WHO. Guideline for the treatment of malaria, 2015:

<http://www.who.int/malaria/publications/atoz/9789241549127/en>

4.1 Therapeutic indications

WHO. Management of severe malaria, a practical handbook, 2013:

http://apps.who.int/iris/bitstream/10665/79317/1/9789241548526_eng.pdf?ua=1

4.2 Posology and method of administration

WHO: Tropical Medicine and International Health Sept 2014, 9 (Suppl. 1), 7-131:

http://onlinelibrary.wiley.com/doi/10.1111/tmi.12313_2/pdf

Hendriksen IC et al. Population pharmacokinetics of intramuscular artesunate in African children with severe malaria: implications for a practical dosing regimen. Clin Pharmacol Ther. 2013 May;93(5):443-50.

<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3630454/pdf/clpt201326a.pdf>

4.4 Special warnings and precautions for use

Information note on delayed haemolytic anaemia following treatment with artesunate

http://www.who.int/malaria/publications/atoz/who_note_haemolytic_anaemia/en/

Rolling T et al. Delayed Hemolysis After Treatment With Parenteral Artesunate in African Children With Severe Malaria—A Double-center Prospective Study. J Infect Dis 2014; 209: 1921-1928

4.6 Pregnancy and lactation

WHO. Management of severe malaria, a practical handbook, 2013:

http://apps.who.int/iris/bitstream/10665/79317/1/9789241548526_eng.pdf?ua=1

McGready R et al. Artemisinin antimalarials in pregnancy: a prospective treatment study of 539 episodes of multidrug-resistant *Plasmodium falciparum*. *Clin Infect Dis* 2001;33:2009-2016.

Centers for Disease Control and Prevention (CDC). Intravenous artesunate for treatment of severe malaria in the United States 2007: Protocol, IND #76,725. 1-28.

Jansen FH et al. Is artesunate or its active metabolite dihydroartemisinin being excreted in the milk of lactating mothers? *Am J Trop Med Hyg*. 2006;75 Suppl. S :158. Abstract.

Olumide SA, Raji Y. 2011. Long-term administration of artesunate induces reproductive toxicity in male rats. *J Reprod Infertil* 12(4): 249-260.

4.8 Undesirable effects

Maude RJ et al. Does artesunate prolong the electrocardiograph QT interval in patients with severe malaria? *Am J Trop Med Hyg* 2009;80:126-32

Rolling T et al. Delayed Hemolysis After Treatment With Parenteral Artesunate in African Children With Severe Malaria—A Double-center Prospective Study. *J Infect Dis* 2014; 209: 1921-1928

Zoller T et al. Intravenous artesunate for severe malaria in travelers, Europe *Emerging Infect Dis* 2011;17:771-777.

4.9 Overdose

Campos S, de la Cerda P, Rivera A Fatal artesunate toxicity in a child *J Ped Inf Dis* 2008;3:69-75

5.1 Pharmacodynamic properties

Hess KM, Goad JA, Argui PM Intravenous artesunate for the treatment of severe malaria *Ann Pharmacotherapy* 2010;44:1250-1258.

Lin AJ, Klayman DL, Milhous WK Antimalarial activity of new water-soluble dihydroartemisinin derivatives *J Med Chem* 1987;30:2147-2150.

Dondorp AM et al. Artesunate versus quinine for treatment of severe *falciparum* malaria: a randomised trial *Lancet* 2005;366:717-725

Jones KL, Donegan S, Lalloo DG. Artesunate versus quinine for treating severe malaria. *Cochrane Database Syst Rev* 2007;4:CD005967-CD005967

Dondorp AM et al. Artesunate vs. quinine in the treatment of severe *falciparum* malaria in African children (AQUAMAT): an open-label randomized trial *Lancet* 2010; 376: 1647–57

5.2 Pharmacokinetic properties

Nealon C et al. Intramuscular bioavailability and clinical efficacy of artesunate in Gabonese children with severe malaria *Antimicrob Agents Chemother* 2002;46:3933-3939.

Ilett KF et al. The pharmacokinetic properties of intramuscular artesunate and rectal dihydroartemisinin in uncomplicated *falciparum* malaria *Br J Clin Pharmacol* 2002;53:23-30.

Batty KT et al. A pharmacokinetic and pharmacodynamic study of intravenous vs oral artesunate in uncomplicated *falciparum* malaria *Br J Clin Pharmacol* 1998;45:123-129.

Hien TT et al. Comparative pharmacokinetics of intramuscular artesunate and artemether in patients with severe *falciparum* malaria *Antimicrob Agents Chemother* 2004;48:4234-4239.

5.3 Preclinical safety data

Efferth T, Kaina B. Toxicity of the antimalarial artemisinin and its derivatives *Critical Reviews in Toxicol* 2010;40:405-421.