

WHO-PQ RECOMMENDED PATIENT INFORMATION LEAFLET

*This patient information leaflet focuses on uses of the medicine covered by WHO's Prequalification Team - Medicines. The recommendations for use are based on WHO guidelines and on information from stringent regulatory authorities.**

The medicine may be authorised for additional or different uses by national medicines regulatory authorities.

* https://extranet.who.int/pqweb/sites/default/files/documents/75%20SRA%20clarification_Feb2017_newtempl.pdf

Information for the patient

[MA102 trade name][†]
Amodiaquine (hydrochloride)/artesunate

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have questions about the medicine, ask your health care provider.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness seem to be the same as yours.
- If you are concerned about any side effects, talk to your health care provider. This includes unwanted effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What [MA102 trade name] is and what it is used for
2. What you need to know before you take [MA102 trade name]
3. How to take [MA102 trade name]
4. Possible side effects
5. How to store [MA102 trade name]
6. Contents of the pack and other information

1. What [MA102 trade name] is and what it is used for

[MA102 trade name] is used to treat malaria. It contains two antimalarial medicines, artesunate and amodiaquine, which work together to kill the malaria parasite in your body. Your health care provider has recommended [MA102 trade name] for you because you have malaria.

It is to be used only for treating uncomplicated malaria caused by *Plasmodium falciparum* (a type of malaria parasite). Uncomplicated malaria is when the patient is conscious and is not suffering serious symptoms such as severe anaemia, jaundice, breathing difficulties and kidney failure. For complete cure it is important that you finish taking all the tablets that your health care provider has prescribed for you.

2. What you need to know before you take [MA102 trade name]

Do not take [MA102 trade name]

- if you are allergic to artesunate, amodiaquine or to any of the other ingredients of [MA102 trade name] (see section 6, What [MA102 trade name] contains);
- if you have ever had liver problems during treatment with amodiaquine;
- if you have ever suffered from sudden high fevers, shaking, severe sore throat or ulcers in the mouth (symptoms of low white blood cell count) during treatment with amodiaquine;
- if you have an eye disease with damage to your retina.

If you have any doubt, check with your health care provider.

Take special care with [MA102 trade name]

This medicine may rarely damage the liver or cause blood problems that can be fatal. If you feel weak, have nausea, vomiting, abdominal pain, jaundice (yellowing of the skin with dark urine, pale bowel motions), sore throat or mouth ulcers, stop treatment with [MA102 trade name] and see your health care provider immediately; you may need blood tests.

[†] Trade names are not prequalified by WHO. This is the national medicines regulatory agency's responsibility.

Tell your health care provider if you have used this or another amodiaquine-containing medicine very recently. This is because you are more likely to get side effects.

This medicine may cause abnormal movements of the head, tongue or neck (extrapyramidal symptoms, see section 4, Possible side effects). These symptoms are usually reversed by stopping treatment. If these rare effects occur, stop the treatment and speak with your health care provider immediately. You may need other treatment for these symptoms. Your doctor may recommend another malaria treatment.

Tell your health care provider if:

- you have ever had a problem with your heart rhythm or any other heart problem;
- a close relative has had heart rhythm problem or died suddenly;
- you have low levels of potassium or magnesium in your blood;
- you have problems with your liver or kidneys.

If in doubt, ask your health care provider.

Taking other medicines

Tell your health care provider if you are taking or have recently taken any other medicines, including medicines obtained without a prescription. Some medicines and [MA102 trade name] interfere with each other. Some of these medicines include:

- other medicines used to treat malaria,
- medicines used to treat an abnormal heart rhythm,
- medicines that can have side effects on your heart, including some medicines used to treat depression or mental illnesses, and some antibiotic medicines,
- medicines used to treat HIV/AIDS.

Pregnancy and breast-feeding

If you are pregnant, you must check with your health care provider before you take this medicine, particularly during the first 3 months of pregnancy.

If you learn that you are pregnant during the treatment, speak with your health care provider.

You can breast-feed your baby during treatment with this medicine but check with your health care provider first.

Driving and using machines

If you feel dizzy or tired after taking this medicine, you should not drive or operate machinery.

[MA102 trade name] contains lactose

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

3. How to take [MA102 trade name]

Always take [MA102 trade name] exactly as your health care provider has told you. You should check with your health care provider if you are not sure.

The dose depends on the patient's weight as follows:

Weight (approximate age)	Dose
4.5 kg to 9 kg (2 to 11 months)	1 tablet [MA102 trade name] daily for 3 days (25 mg artesunate + 67.5 mg amodiaquine daily for 3 days)
9 kg to 18 kg (1 to 5 years)	2 tablets [MA102 trade name] daily for 3 days (50 mg artesunate + 135 mg amodiaquine daily for 3 days)
18 kg to 36 kg (6 to 13 years)	4 tablets [MA102 trade name] daily for 3 days (100 mg artesunate + 270 mg amodiaquine daily for 3 days)

more than 36 kg (over 14 years)	8 tablets [MA102 trade name] daily for 3 days (200 mg artesunate + 540 mg amodiaquine daily for 3 days)
------------------------------------	--

Do not take [MA102 trade name] with a meal that contains a high amount of fat.

Swallow the tablet with a drink of water.

For very young children or patients who cannot swallow the tablets whole, the tablets can be crushed and swallowed with water, or mixed with water and swallowing the mixture.

If you vomit within half-an-hour of taking the tablets, take another dose as soon as you can. Then contact your health care provider for more tablets to finish the course. Tell your health care provider if you keep vomiting.

If you take more [MA102 trade name] than you should

If you accidentally take too many tablets, then you or somebody else should immediately contact a health care provider or the nearest hospital emergency department. An overdose of this medicine can cause headaches, dizziness, visual disturbances, a drop in blood pressure, convulsions (fits), or abnormal movements of the head, tongue or neck. If any of these symptoms occur, see a health care provider immediately.

If you forget to take [MA102 trade name]

Make sure that you do not miss a dose of this medicine. However, if you forget a dose, take the missed dose as soon as you remember. Take the next dose as usual. Do not take a double dose to make up for a forgotten tablet.

If you stop taking [MA102 trade name]

For this medicine to work properly you must take it regularly until you finish the course. Even if you feel better, it does not mean that your malaria has been completely cured. Tiredness, nausea and vomiting may be caused by the infection and not by the medicine so you should not stop your medicine if you get these effects.

If you have any further questions on the use of this medicine, ask your health care provider.

4. Possible side effects

Like all medicines, [MA102 trade name] can cause side effects, although not everybody gets them.

Common side effects (affects 1 to 10 users in 100): loss of appetite, difficulty sleeping, sleepiness, cough, nausea, abdominal pain, weakness, tiredness.

Uncommon side effects (affects 1 to 10 users in 1,000): acute bronchitis, gastroenteritis, oral thrush, decrease in red blood cell count (anaemia), hypoglycaemia (low blood sugar), hallucination, tingling and numbness of the limbs, yellowing of the eye, dizziness, heart rhythm disorders, diarrhoea, vomiting, itching, rashes, swelling of the face, skin disorders, joint pain, swelling of the limbs (oedema), fever.

Side effects that may be linked to amodiaquine, one of the active ingredients of [MA102 trade name]: low number of white blood cells (which can make you more prone to infections), potentially severe disturbances of liver function, eye disorders, slate-grey pigmentation, especially of the fingers and mucous membranes (lining of body cavities), nervous system problems and muscle impairment.

During post-marketing experience the following additional side effects have been reported: Common side effects (affects 1 to 10 users in 100): alteration of the heart rhythm (called 'prolongation of QT interval').

Frequency not known: headache, cold, flu, rhinitis (runny nose), shivering, sore throat, swelling of the spleen, convulsion (fits), yellowing of the skin, allergic reaction. Some of these effects may be caused by the malaria.

Also, side effects called extrapyramidal symptoms can occur. These symptoms include abnormal movements of the head, tongue or neck (e.g. facial spasms, tongue protrusion, difficulties in speaking) and rigidity in the body, even after a single dose.

If any of the side effects gets serious, or if you notice any side effect not listed in this leaflet, tell your health care provider.

Reporting of side effects

If you get any side effects, talk to your health care provider. This includes unwanted effects not listed in this leaflet. If available, you can also report side effects directly through the national reporting system. By reporting side effects you can help improve understanding about the safety of this medicine.

5. How to store [MA102 trade name]

Do not store above 30°C.

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date stated on the blister and carton after EXP. The expiry date refers to the last day of that month.

Do not throw away any medicines in wastewater or household waste. Ask your health care provider how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What [MA102 trade name] contains

- The active ingredients are 25 mg artesunate and 67.5 mg amodiaquine (as hydrochloride).
- The other ingredients of [MA102 trade name] are: lactose anhydrous, lactose monohydrate, croscarmellose sodium, magnesium stearate, calcium carbonate DC 95S with 5% corn starch and colloidal anhydrous silica.

What [MA102 trade name] looks like and contents of the pack

[MA102 trade name] is a circular uncoated bilayered flat bevelled tablet with a white to light yellowish layer with a central break-line and a yellow layer debossed with the figure '25'.

The break-line is only to facilitate breaking for ease of swallowing and not to divide into equal doses.

The tablets are provided in an Alu/Alu blister pack containing 3 tablets. A carton contains 1 or 25 such blisters.

Supplier and Manufacturer

Supplier

Cipla Limited
Cipla House,
Peninsula Business Park,
Ganpatrao Kadam Marg,
Lower Parel, Mumbai- 400013,
India
Phone: 91-22-24826000
Fax: 91-22-24826120
E-mail: globalra@cipla.com

Manufacturer

Cipla Limited
Unit II, A-42, MIDC
Patalganga, Raigad District
Maharashtra, 410 220,
India

For any information about this medicine, contact the local representative of the supplier:

This leaflet was last revised in September 2021

Detailed information on this medicine is available on the World Health Organization (WHO) website: <https://extranet.who.int/pqweb/medicines>