

WHO-PQ RECOMMENDED PATIENT INFORMATION LEAFLET

*This patient information leaflet focuses on uses of the medicine covered by WHO's Prequalification Team - Medicines. The recommendations for use are based on WHO guidelines and on information from stringent regulatory authorities.**

The medicine may be authorised for additional or different uses by national medicines regulatory authorities.

* https://extranet.who.int/pqweb/sites/default/files/documents/75%20SRA%20clarification_Feb2017_newtempl.pdf

[#] Formerly known as 'Strides Shasun Limited'.

Information for the patient

[MA110 trade name][†]
Artemether/lumefantrine

*The warnings and instructions in this leaflet are intended for the person taking the medicine.
If you are a parent or carer responsible for giving the medicine to someone else such as a child,
you will need to apply the instructions accordingly.*

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have questions about the medicine, ask your health care provider.
- This medicine is for you only. Do not pass it on to others. It may harm them, even if their illness seems to be the same as yours.
- If you are concerned about any side effects, talk to your health care provider. This includes unwanted effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What [MA110 trade name] is and what it is used for
2. What you need to know before you take [MA110 trade name]
3. How to take [MA110 trade name]
4. Possible side effects
5. How to store [MA110 trade name]
6. Contents of the pack and other information

1. What [MA110 trade name] is and what it is used for

[MA110 trade name] is a medicine used to treat malaria in adults and children. It contains two active substances, artemether and lumefantrine, which work together to kill the malaria parasite.

Malaria is caused by infection with a parasite called *Plasmodium*, spread by the bite of an infected mosquito. [MA110 trade name] is used when the malaria is caused by a type of malaria parasite called *Plasmodium falciparum* and only when the infection is not severe enough to affect the brain or other key organs.

Your health care provider will follow the most recent official guidelines on the use of malaria medicines to select the right medicine for your malaria treatment.

2. What you need to know before you take [MA110 trade name]

Do not take [MA110 trade name]:

- if you are allergic (hypersensitive) to artemether, lumefantrine, or any of the other ingredients of this medicine (listed at the end of this leaflet). If you think you may be allergic, ask your health care provider for advice.
- if your health care provider tells you that you have severe malaria (affecting the brain or with complications affecting other organs such as lungs or kidneys).
- if you have a heart condition, such as dangerously irregular heartbeats (arrhythmias), an alteration in the electrical activity of the heart called prolonged QT interval, slow heartbeat, or severe heart disease.

[†] Trade names are not prequalified by WHO. This is the national medicines regulatory agency's responsibility.

- if any member of your family (e.g. parents, grandparents, brothers and sisters) has died suddenly due to a heart rate problem or is known to have been born with heart rate problems.
- if you are taking certain medicines (see “Other medicines and [MA110 trade name]”).
- if you have low blood levels of electrolytes such as potassium or magnesium.

If any of these apply to you, **tell your health care provider before taking [MA110 trade name]**.

Warnings and precautions

Talk to your health care provider **before taking [MA110 trade name]**:

- If you have severe liver or kidney problems. Your health care provider may recommend extra tests to monitor your heart and the level of potassium in your blood.

Take special care with [MA110 trade name]:

- If your condition worsens, or if you feel too unwell to eat and drink, contact your health care provider immediately. Your health care provider may want to perform a test called an electrocardiogram (ECG) and check the levels of electrolytes, such as potassium and magnesium in your blood before and during treatment.
- If you are taking or have taken any other medication for the treatment of malaria, talk to your health care provider, because some of these medicines must not be given together with [MA110 trade name].
- [MA110 trade name] is used to treat malaria due to *Plasmodium falciparum*. If you are also infected with another type of malaria parasite, *Plasmodium vivax*, your health care provider will give you another medicine for you to take after completing [MA110 trade name] treatment.

Other medicines and [MA110 trade name]

It is important that you tell your health care provider if you are taking or have recently taken any other medicines, including medicines obtained without a prescription. These may affect the action of [MA110 trade name], or [MA110 trade name] may affect their action. Side effects of either medicine may become worse and/or the medicines may become less effective.

Especially tell your health care provider if you take or have recently taken:

- Any other medicines to treat or prevent malaria
- Medicines for your heart
- Antipsychotic medicines (for treatment of abnormal condition of the mind)
- Antidepressants (medicines for depression and low mood)
- Antibiotics
- Antihistamines (for treatment of, e.g., allergies)
- Medicines to treat HIV infection
- Medicines to treat hepatitis B or hepatitis C infection
- Medicines against fungal infection
- Hormonal methods of birth control (for example birth control pills or contraceptive patch)

[MA110 trade name] with food and drink

[MA110 trade name] should be taken with food or a milky drink.

Pregnancy, breast-feeding and fertility

Pregnancy

[MA110 trade name] can be used during pregnancy.

Breast-feeding

The amounts of the active substances of [MA110 trade name] that pass into breast milk are low and [MA110 trade name] can be used during breast-feeding.

Fertility

There is no information on the effects of [MA110 trade name] on fertility in humans.

Driving and using machines

[MA110 trade name] may cause dizziness and tiredness. If you feel dizzy or tired while taking [MA110 trade name], do not drive and do not use any tools or machines.

3. How to take [MA110 trade name]

Your health care provider will explain to you how many tablets of [MA110 trade name] to take. The dose depends on your weight. You should take the medicine for 3 days, with no breaks in between.

The usual doses of the medicine for patients of different weights are described below:

Patient's weight	Time					
	Day 1		Day 2		Day 3	
	<i>Immediately after diagnosis/onset of symptoms</i>	<i>8 hours after previous dose</i>	<i>12 hours after previous dose</i>	<i>12 hours after previous dose</i>	<i>12 hours after previous dose</i>	<i>12 hours after previous dose</i>
From 5 kg up to 15 kg	1 tablet	1 tablet	1 tablet	1 tablet	1 tablet	1 tablet
From 15 kg up to 25 kg	2 tablets	2 tablets	2 tablets	2 tablets	2 tablets	2 tablets
From 25 kg up to 35 kg	3 tablets	3 tablets	3 tablets	3 tablets	3 tablets	3 tablets
From 35 kg	4 tablets	4 tablets	4 tablets	4 tablets	4 tablets	4 tablets

Disperse the tablets in drinking water before taking the dose, as described under 'Instructions for taking [MA110 trade name] tablets', below.

Take the first dose immediately when your health care provider has diagnosed malaria.

Take the second dose 8 hours after the first dose.

Then take the remaining doses 12 hours apart.

If you vomit within 1 hour of taking the medication, you should repeat the dose.

You should always take [MA110 trade name] exactly as described by the health care provider and it is important to complete the course of tablets that is recommended. Check with your health care provider if you are not sure.

Instructions for taking [MA110 trade name] tablets

[MA110 trade name] should be taken with food or a milky drink. If you are unable to tolerate food, [MA110 trade name] should still be taken, but your body may take up less of the medicine.

Each tablet should be dispersed in a minimum of 10 mL water; the maximum volume of water recommended for dispersion of a dose is 50 mL.

1. Look up the patient's weight in the table above.
2. Then look for the number of tablets that will be required.
3. Place the required volume of water in a small, clean cup or bowl and add the required number of tablets.
4. Stir gently until the tablets disperse.
5. Drink the entire medicine mixture immediately.

6. Rinse the cup with a little additional water and drink this water.

If you take more [MA110 trade name] than you should

If you take too many tablets, immediately contact your health care provider or the nearest hospital emergency department for further advice. Bring the medicine box with you, so that you can easily describe what you have taken.

If you forget to take [MA110 trade name]

Try to make sure that you do not miss any dose. However, if you do forget a dose, take the missed dose as soon as you realise that you have forgotten it. Then take the next dose after the prescribed interval. Do not take a double dose to make up for a forgotten tablet. **Make sure you take all six doses of this regimen.**

If you stop taking [MA110 trade name]

You should keep taking the medicine for as long as your health care provider has ordered, even if you are feeling better. If you stop the medicine too soon, the infection may not be completely cured.

If you have any further questions on the use of this product, ask your health care provider.

4. Possible side effects

Like all medicines, [MA110 trade name] can cause side effects, although not everybody gets them. It is important that you inform the health care provider of any change in your health.

Most of the side effects are mild to moderate and generally disappear after a few days to a few weeks after treatment.

However, some side effects could be serious and need immediate medical attention. Allergic (hypersensitivity) reactions may occur rarely, in up to 1 person in 1000.

- If you get a rash, swelling of the face, lips, tongue or throat with difficulty in swallowing or breathing, **tell your health care provider straight away.**

Of the other side effects that may occur with [MA110 trade name], the *most commonly* reported side effects (in more than 1 in every 10 patients treated) include

- fast heart beat (palpitations)
- headache
- dizziness
- cough
- nausea (feeling sick) or vomiting (being sick)
- abdominal (belly) pain
- decreased appetite
- joint pain
- muscle pain
- general weakness or tiredness
- sleep disorders.

Other *common* side effects (reported in more than 1 in every 100 patients treated) include

- changes in the electrical activity of the heart (QT-prolongation) seen in tests
- abnormal blood tests for liver function
- rash
- itching
- diarrhoea
- abnormal walking*
- needles and pins (paraesthesia) or numbness of the hands and feet*
- involuntary, rhythmic, muscular contractions (clonus)
- insomnia.

Uncommon side effects (greater than 1 in every 1000 patients treated but less than 1 in 100) include

- clumsiness or difficulty in making movements smoothly (ataxia)*
- decreased skin sensitivity*
- sleepiness
- nettle rash (urticaria).

*These side effects have been reported in adults and adolescents above 12 years of age.

There have also been some reports of anaemia (low numbers of red blood cells) due to breakdown of the red blood cells, which has been reported up to a few weeks after treatment has been stopped (delayed haemolytic anaemia). It is not clear how often this may occur.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your health care provider as soon as possible.

Reporting of side effects

If you get a side effect, talk to your health care provider. This includes side effects not listed in this leaflet. You may also be able to report such effects directly to your national reporting system if one is available. By reporting side effects, you can help to improve the available information on this medicine.

5. How to store [MA110 trade name]

Store below 30°C. Avoid excursions above 30°C. Protect from light and moisture.

Store tablets in blisters in the provided carton.

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date stated on the carton. The expiry date refers to the last day of that month.

Do not use this medicine if you notice that it is different from the description below.

Do not throw away any medicines in wastewater or household waste. Ask your health care provider how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What [MA110 trade name] contains

- The active ingredients are artemether and lumefantrine. Each tablet contains 20 mg artemether and 120 mg lumefantrine.
- The other ingredients of [MA110 trade name] are: microcrystalline cellulose, croscarmellose sodium, hypromellose, polysorbate 80, colloidal silicon dioxide, crospovidone, magnesium stearate, saccharin sodium, cherry flavour.

There is too little sodium in this medicine to have any effect, even if you are on a low-sodium diet.

What [MA110 trade name] looks like and contents of the pack

[MA110 trade name] is a yellow coloured, circular, flat bevel-edged tablet with score line on one side and plain on the other side.

The score line is not intended for breaking the tablet.

[MA110 trade name] is packed in PVC/PE/PVDC-Alu blister. Each blister contains either 6 or 12 tablets. One such blister is packed in a mono carton. 30 such mono cartons are packed in an outer carton.

Supplier and Manufacturer

Supplier

Strides Pharma Global PTE Limited
1 Gateway Drive
#06-06
Westgate Tower

Manufacturer

Strides Pharma Science Limited
KRS Gardens
Tablet Block
36/7, Suragajakkanahalli

Singapore 608531
Tel. No.: +(65) 6421 4680
Email: AshokKumar.J@strides.com,
info@strides.com

Indlavadi cross
Aanekal Taluk
Bangalore 562106
India
Tel. No.:91-80-67840600
Fax No: 91-80-67840606
Email: AshokKumar.J@strides.com,
info@strides.com

Universal Corporation Limited
Club Road
Past Kikuyu Post Office
Plot No. 13777
P.O Box: 1748 - 00902
Kikuyu Town
Kikuyu
Kenya

For any information about this medicine, contact the local representative of the supplier:

Region: **East Africa**
Country: **Kenya**
Address: **M/s. Universal Corporation Ltd,**
Club Road, Past Kikuyu Post Office
P.O Box 1748-00902
Kikuyu, Kenya
Contact Person: Mr. Palu Dhanani
Email Id: palu.dhanani@ucl.co.ke

Region: **East Africa**
Country: **Uganda**
Address: **M/s. Delmaw Enterprises Limited**
Plot 9/5 & 6 , Gathani House
Bombo Road
P.O Box No. 1574
Kampala – Uganda
Contact Person: Mr. Gerald
Email Id: delmaw.regulatory@artemislife.com

Region: **East Africa**
Country: **Tanzania**
Address: **M/s. Samiro Pharmaceuticals Ltd**
P.O.BOX 38062
Uhuru Street, Kariakoo,
Dar-es-Salaam, Tanzania
Contact Person: Mr. Abbas Mohammed
Email Id: info@samiropharma.com

Region: **Central Africa**
Country: **DR Congo**
Address: **M/s. Gulf Africa investments Sarl**
(Division: Gulf Africa Pharma)
B.P. 14009
Avenue du Marche No: PL 6356
Commune de la Gombe

Artemether/lumefantrine 20mg/120mg
dispersible tablets (Strides Pharma
Global PTE Ltd)[#], MA110

WHOPAR Part 3

March 2023
Section 6 updated : May 2024

Kinshasa RD- Congo

Contact Person: Mr. Leonard Netto Email Id: leonard.netto@artemislife.com

Region: **West Africa**

Country: **Nigeria**

Address: **M/s. Strides Lifesciences Ltd,**

Chenline House

7,Obasa St. Off Oba Akran Ave

Ikeja, Lagos, Nigeria

Contact Person: Mr. Smatt Aguba

Email Id: SP@strides.com

Region: **Southern Africa**

Country: **Zambia**

Address: **M/s. Yash Pharmaceuticals Ltd**

Plot 27 Lilayi Road, Lusaka

P.O BOX: 34562 Zambia

Contact Person: Mr. Himanshu Patel

Email Id: h.patel@yashpharmaceuticals.com

This leaflet was drafted in March 2023

Section 6 was updated in May 2024

Detailed information on this medicine is available on the World Health Organization (WHO) website:
<https://extranet.who.int/pqweb/medicines>