

WHO-PQ RECOMMENDED PATIENT INFORMATION LEAFLET

*This patient information leaflet focuses on uses of the medicine covered by WHO's Prequalification Team - Medicines. The recommendations for use are based on WHO guidelines and on information from stringent regulatory authorities.**

The medicine may be authorised for additional or different uses by national medicines regulatory authorities.

Information for the patient

[MA141 trade name][†]

Dihydroartemisinin /piperaquine phosphate 20 mg/160 mg dispersible tablets

If you are a carer or parent looking after the person who takes this medicine, use this leaflet to give the medicine correctly and take note of the warnings and side effects.

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have questions about the medicine, ask your health care provider.
- This medicine is for you only. Do not pass it on to others. It may harm them, even if their illness seems to be the same as yours.
- If you are concerned about any side effects, talk to your health care provider. This includes unwanted effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What [MA141 trade name] is and what it is used for
2. What you need to know before you take [MA141 trade name]
3. How to take [MA141 trade name]
4. Possible side effects
5. How to store [MA141 trade name]
6. Contents of the pack and other information

1. What [MA141 trade name] is and what it is used for

[MA141 trade name] is a medicine used to treat uncomplicated malaria (when the infection is not severe enough to affect the brain or other key organs). It contains the active substances dihydroartemisinin and piperaquine phosphate, which work together to kill the parasites that cause malaria.

Malaria is caused by infection with a parasite called *Plasmodium*, spread by the bite of an infected mosquito. There are different types of *Plasmodium* parasite. [MA141 trade name] kills all types of *Plasmodium* parasite.

Your health care provider will follow the most recent official guidelines on the use of malaria medicines to select the right medicine for your malaria treatment.

2. What you need to know before you take [MA141 trade name]

Do not take [MA141 trade name] if you:

- are allergic (hypersensitive) to the active substances, dihydroartemisinin or piperaquine phosphate, or to any of the other ingredients of [MA141 trade name] (see section 6);
- have a severe type of malaria infection which has affected parts of the body such as the brain, lungs or kidneys (your health care provider will check this before giving the medicine);
- have a heart condition affecting the rhythm or electrical activity of the heart, or have an unusually slow heart rate;
- know that any member of your family (parents, grandparents, brothers or sisters) died suddenly due to a heart problem or was born with heart problems;

[†] Trade names are not prequalified by WHO. This is the national medicines regulatory agency's responsibility.

- suffer from changes to the levels of salts in the body (electrolyte imbalances);
- are taking other medicines that can have an effect on heart rhythm, such as:
 - the heart medicines quinidine, disopyramide, procainamide, amiodarone, dofetilide, ibutilide, hydroquinidine or sotalol;
 - medicines used to treat depression;
 - medicines used to treat mental health problems such as phenothiazines, sertindole, sultopride, chlorpromazine, haloperidol, mesoridazine, pimozide, or thioridazine;
 - medicines used to treat infections. These include some of the types of medicines used to treat bacterial infections (macrolides [such as erythromycin or clarithromycin] and fluoroquinolones [such as moxifloxacin and sparfloxacin]) or fungal infections (including fluconazole and imidazole) as well as pentamidine (used to treat a specific type of pneumonia) and saquinavir (for treatment of HIV);
 - antihistamines used to treat allergies or inflammation such as terfenadine, astemizole or mizolastine;
 - certain medicines used to treat stomach problems such as domperidone or droperidol;
 - other medicines such as vinca alkaloids and arsenic trioxide (used to treat certain cancers), bepridil (used to treat angina), diphemanil (used to treat stomach disturbances), levomethadyl and methadone (used to treat drug addiction), and probucol (used to treat high blood cholesterol levels).
- have recently (for example within about one month) been treated for malaria with certain medicines or taken certain medicines to prevent malaria. These medicines include: mefloquine, halofantrine, lumefantrine, chloroquine or quinine.

If any of the above applies or if you are unsure, tell your health care provider before taking [MA141 trade name].

Take special care with [MA141 trade name]

Check with your health care provider before taking this medicine if you (or a child taking the medicine):

- have liver or kidney problems.
- are taking or have taken any other medicines for the treatment of malaria (other than those mentioned above), especially if these were given by injection or drip.
- are pregnant or breastfeeding (see below).
- are elderly (over 65 years)
- are vomiting (especially in young children).
- are taking any other medicines which could interact with [MA141 trade name]. Examples are listed in the section “Taking other medicines”.

If you are not sure about any of the above, please ask your health care provider.

Taking other medicines

Please tell your health care provider if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

Other medicines can affect the way [MA141 trade name] works or increase the risk of side effects if taken together, so your health care provider may decide that [MA141 trade name] is not suitable or that extra checks are needed.

As well as medicines that affect the heart rhythm and some malaria medicines (see ‘**Do not take [MA141 trade name]**’, above) other medicines that may interact with [MA141 trade name] include:

- some *medicines for HIV*
- medicines to treat *infections such as tuberculosis*
- medicines for *epilepsy*
- *hormonal contraceptives*
- some *medicines for heart problems, high blood pressure or high cholesterol*

- medicines such as *ciclosporin* (used after organ transplants or for autoimmune disease), *midazolam* (for anxiety or sleep problems), *nefazodone* (used to treat depression), *omeprazole* (for stomach acid), *paracetamol* (a painkiller), and *theophylline* (to help breathing)
- some herbal medicines such as *St John's wort* (for anxiety).

Make sure you tell your health care provider about any medicines you may take.

Taking [MA141 trade name] without food and drink

You should take [MA141 trade name] tablets with water only.

You should take this medicine on an empty stomach as food, especially fatty or oily food, can increase the risk of side effects. You should take each dose at least 3 hours after the last time you ate, and no food should be taken within 3 hours after each dose of [MA141 trade name]. For example, if you eat at midday, you can take your dose after 3 pm and then eat again after 6 pm, or if you eat at 6 pm, you can take your dose at 9 pm and then eat your breakfast at your normal time the next morning.

You can drink water at any time. You should not take [MA141 trade name] with grapefruit juice due to possible interactions.

Pregnancy and breast-feeding

Tell your health care provider if you are pregnant, think you may be pregnant or become pregnant, or if you are breast-feeding.

[MA141 trade name] is one of several malaria treatments that can be used in pregnancy – your health care provider will select the most appropriate malaria medicine for your situation. If you are given [MA141 trade name] while pregnant, your health care provider may carry out extra checks on how your pregnancy progresses.

You may breast-feed your baby while taking this medicine.

If you are taking folate supplements to reduce the risk of birth defects, you can continue taking them at the same time as [MA141 trade name].

Ask your health care provider for advice before taking any medicine during pregnancy or breast-feeding.

Driving and using machines

[MA141 trade name] is not likely to affect your ability to drive or operate machinery.

However, make sure you feel well enough to take on any skilled tasks.

3. How to take [MA141 trade name]

Always take [MA141 trade name] exactly as your health care provider has told you to. You should check with your health care provider if you are not sure.

Your health care provider will explain to you how many tablets of [MA141 trade name] to take. The dose depends on the patient's weight.

You should take the medicine for 3 days, with no breaks in between. You should try to take the dose at about the same time on each of the 3 days.

The usual doses of the medicine for patients of different weights are described below:

Body weight	Dose
5 kg to less than 8 kg	1 tablet per day
8 kg to less than 11 kg	<i>different tablet needed</i>
11 kg to less than 17 kg	2 tablets per day
17 kg to less than 25 kg	3 tablets per day

Children weighing **8 to less than 11 kg**, and **patients weighing 25 kg** or more should be given a different tablet containing a more suitable dose. Tell your health care provider if this applies so that they can prescribe a suitable tablet.

Disperse the tablets in drinking water before taking the dose. Each tablet should be dispersed in a minimum of 5 to 10 mL water; the maximum volume of water recommended for dispersion of a dose is 50 mL.

You should take each dose at least 3 hours after your last meal. You should also avoid eating until at least 3 hours after taking [MA141 trade name]. For example, if you eat at midday, you can take your dose at 3 pm and then eat again after 6 pm, or if you eat at 6 pm, you can take your dose at 9 pm and then eat your breakfast at your normal time the next morning. This is to ensure the stomach is empty when taking the medicine, as food in the stomach, especially fatty or oily food, can increase the risk of side effects. Drinking water is permitted at any time.

Taking [MA141 trade name]

1. Look up your weight in the table above.
2. Then look for the number of tablets that will be required.
3. Place the required volume of water in a small, clean cup or bowl and add the required number of tablets.
4. Stir gently until the tablets disperse.
5. Take the whole mixture to drink right away.
6. Rinse the cup with a little additional water and drink this water too, to make sure all of the dose gets swallowed.
7. Repeat this procedure on each of the 3 days of the course.

Vomiting when taking this medicine:

If you vomit within 30 minutes after taking [MA141 trade name], the **whole dose** must be taken again.

If you vomit between half an hour and 1 hour after the dose, **half the dose** must be taken again. If you vomit after this time, you do not need to take another dose.

If you also vomit after the second dose, do not take another dose. Contact your health care provider urgently to obtain an alternative treatment for malaria.

If you take more [MA141 trade name] than you should:

If you take more than the recommended dose, tell your health care provider. Your health care provider may suggest special monitoring for you because doses higher than those recommended may have an unwanted, severe effect on the heart (see also section 4).

If you forget a dose of [MA141 trade name]:

If you forget to take the second dose of [MA141 trade name] at the right time, take it as soon as you remember. Then take the third (last) dose approximately 24 hours after the second dose.

If you forget to take the third (last) dose at the right time, take it as soon as you remember.

Never take more than one dose on the same day to make up for a missed dose. Check with your health care provider if you are not sure.

If your stop taking [MA141 trade name]:

For the medicine to work effectively, you should take the tablets as instructed and should complete the 3 days course of treatment. If you are not able to do this, talk to your health care provider.

Taking this medicine, if the malaria infection returns:

If you get another attack of malaria within a year of taking [MA141 trade name], your health care provider may decide that a second course of this medicine can be taken. However, patients must not take more than two courses within one year.

If you get malaria more often than this or get malaria again within a month or two after taking [MA141 trade name], talk to your health care provider. You will be prescribed another treatment.

If you have any further questions on the use of this medicine, ask your health care provider.

4. Possible side effects

Like all medicines, [MA141 trade name] can cause side effects, although not everybody gets them. Most of the side effects are not severe and normally disappear within a few days or weeks after treatment.

If you get a rash, swelling of the face, lips, tongue or throat with difficulty in swallowing or breathing, these may be signs of an allergic reaction. Tell your health care provider immediately or go immediately to the emergency department of your nearest hospital, taking this leaflet with you.

A heart problem called QT prolongation can occur in some people who take [MA141 trade name], particularly those who have a heart condition or are taking another medication that can cause the same problem. Tell your health care provider if you have a heart condition or are taking any medications. Your health care provider can advise you about using [MA141 trade name].

If you notice anything different about your heart rhythm or have symptoms (such as palpitations or irregular heartbeat) you should contact your health care provider as soon as possible and before the next dose is due.

Problems with red blood cells

Sometimes damage to your red blood cells, called haemolytic anaemia can occur after receiving malaria treatment, especially in young children and if you have previously received artesunate injections to treat malaria. This can happen up to one month following treatment with [MA141 trade name]. In most cases, the anaemia recovers without specific treatment, but if it is severe a blood transfusion may be required. If the breakdown of red blood cells is suspected to be caused by your immune system (autoimmune haemolytic anaemia) you may need further treatment. Your health care provider will carry out tests to check your blood.

If you get one or more of the following symptoms after treatment with [MA141 trade name] you should contact your health care provider immediately: pale skin, general weakness, headache, shortness of breath and rapid heartbeat; particularly with exercise, confusion, dizziness, or dark-coloured urine.

Side effects in adults:

Common (affecting less than 1 in 10 patients but more than 1 in 100):

Low red blood cell counts (anaemia), headache, heart rhythm disturbances (ECG changes or noticing unusually fast heart beats or palpitations), fever, general weakness.

Uncommon (affecting less than 1 in 100 patients but more than 1 in 1000):

Flu, respiratory infections, poor appetite or loss of appetite, dizziness, convulsions (fits), irregular or slow heart rate, cough, vomiting, abdominal pain, diarrhoea, nausea, inflammation or enlargement of the liver, abnormal liver function tests, damage to liver cells, itching, pain in the muscles or joints.

Side effects in children:

Very common (affecting more than 1 in 10 patients):

Flu, cough, fever.

Common (affecting less than 1 in 10 patients but more than 1 in 100):

Respiratory infections, ear infection, low red blood cell counts (anaemia), abnormalities in various types of blood cells (white blood cells and platelets), poor appetite or loss of appetite, eye infection, heart rhythm disturbances (changes as in adults, ECG changes), abdominal pain, vomiting, diarrhoea, skin inflammation, rash, general weakness.

Uncommon (affecting less than 1 in 100 patients but more than 1 in 1000):

Abnormalities in red blood cells, excessive numbers of platelets, enlargement of some organs (such as liver or spleen), swollen lymph glands, convulsions (fits), headache, abnormal heart sounds (heard by your doctor with a stethoscope), nose bleeds, runny nose, nausea, inflammation of the mouth, inflammation or enlargement of the liver, yellowing of the skin and eyes, abnormal liver function blood tests, skin itching and inflammation, pain in the joints.

Reporting of side effects

If you get a side effect, talk to your health care provider. This includes side effects not listed in this leaflet. You may also be able to report such effects directly to your national reporting system if one is available. By reporting side effects, you can help to improve the available information on this medicine.

5. How to store [MA141 trade name]

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date stated on the label after EXP.

Do not use this medicine if you notice description of the visible signs of deterioration that it is different from the description below.

Do not throw away any medicines in wastewater or household waste. Ask your health care provider how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What [MA141 trade name] contains

- The active ingredients are dihydroartemisinin and piperazine phosphate.
- The other ingredients of [MA141 trade name] are pregelatinized starch, microcrystalline cellulose, dextrin, croscarmellose sodium, magnesium stearate, sweet flavour, orange flavour and sucralose.

There is too little sodium in this medicine to have any effect, even if you are on a low-sodium diet.

What [MA141 trade name] looks like and contents of the pack

[MA141 trade name] are off-white or pale yellow, round, uncoated tablets. They are flat on the top and bottom with flat edges. The tablets have 'D' debossed (stamped into) one side and a break line on the other side.

The break line is only to facilitate breaking for ease of swallowing and not to divide the tablet into equal doses.

Nature and contents of the pack

Aluminium foil (PA/Alu/PVC) on aluminium foil blister cards, each containing 3 tablets. Available in boxes of 2 x 3 or 25 x 3 tablets.

Supplier and Manufacturer

Supplier

Guilin Pharmaceutical Co Ltd
No. 43, Qilidian Road, Guilin – 541004
Guangxi
China
Tel.: +86 773 3675053
Fax: +86 773 3675692
E-mail: ra@guilinpharma.com

Manufacturer

Guilin Pharmaceutical Co Ltd
Oral Solid Dosage Workshop 1
No. 43, Qilidian Road, Guilin – 541004
Guangxi
China
Tel.: +86 773 3675053
Fax: +86 773 3675692
E-mail: ra@guilinpharma.com

For any information about this medicine, contact the supplier.

This leaflet was last revised in February 2026

Detailed information on this medicine is available on the World Health Organization (WHO) website:
<https://extranet.who.int/prequal/medicines/prequalified/finished-pharmaceutical-products>