

WHO-PQ RECOMMENDED PATIENT INFORMATION LEAFLET

*This patient information leaflet focuses on uses of the medicine covered by WHO's Prequalification Team - Medicines. The recommendations for use are based on WHO guidelines and on information from stringent regulatory authorities.**

The medicine may be authorised for additional or different uses by national medicines regulatory authorities.

* https://extranet.who.int/prequal/sites/default/files/document_files/75%20SRA%20clarification_Feb2017_newtempl.pdf

Information for the patient

[NT005 trade name][†]
Albendazole

If you are a carer or parent looking after the person who takes this medicine, use this leaflet to give the medicine correctly and take note of the warnings and side effects.

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have questions about the medicine, ask your health care provider.
- This medicine is for you only. Do not pass it on to others. It may harm them, even if their illness seems to be the same as yours.
- If you are concerned about any side effects, talk to your health care provider. This includes unwanted effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What [NT005 trade name] is and what it is used for
2. What you need to know before you take [NT005 trade name]
3. How to take [NT005 trade name]
4. Possible side effects
5. How to store [NT005 trade name]
6. Contents of the pack and other information

1. What [NT005 trade name] is and what it is used for

[NT005 trade name] contains the active substance albendazole. It is used to treat infections caused by some parasites (worms) that usually infect the gut but that can also cause problems in other parts of the body. It works by stopping the worm digesting and using food, so that it starves and dies.

Infections caused by tapeworms

[NT005 trade name] is used to treat *cystic echinococcosis* and *alveolar echinococcosis*, caused by tapeworms that form cysts in the liver, lungs and other parts of the body. The medicine can be used before or after surgery, or when surgery is not possible.

[NT005 trade name] is also used to treat *neurocysticercosis*, an infection of the nervous system caused by pork tapeworms.

[NT005 trade name] may sometimes be given in a community programme to stop the spread of tapeworm infections in the gut (*taeniasis*) in people living in areas where the infection is common.

Lymphatic filariasis

[NT005 trade name] is used to stop the spread of the roundworm infection called *lymphatic filariasis* which can cause fever, cough and wheezing, swelling of the lymph nodes, swollen limbs or genitals, and kidney problems. [NT005 trade name] is used together with other medicines as part of a community programme.

[†] Trade names are not prequalified by WHO. This is the national medicines regulatory agency's responsibility.

Other infections caused by roundworms

[NT005 trade name] is effective against several infections caused by roundworms. It can therefore be used, together with other medicines, as part of a community programme to control the spread of the soil-transmitted infections *ascariasis*, *trichuriasis* and *hookworm* (*ancylostomiasis* and *necatoriasis*).

2. What you need to know before you take [NT005 trade name]

Do not take [NT005 trade name] if you are allergic to albendazole or related medicines such as mebendazole, or any of the other ingredients of this medicine (listed in section 6).

If you are not sure, talk to your health care provider before taking [NT005 trade name].

The warnings and precautions below apply particularly if you are taking a course of treatment with this medicine that lasts several days or longer.

Warnings and precautions

Before starting treatment with [NT005 trade name], tell your health care provider about any medical conditions you have. In particular:

- If you have **liver problems**, you must get medical advice before taking this medicine.

[NT005 trade name] may cause liver problems. Speak with your health care provider straightaway if you are taking a course of treatment with this medicine and you:

- lose your appetite, or get nausea (feel sick), or feel tired all the time, or you have a rash

and you have any of the following

- belly pain, usually on the right side and near the chest
- unexplained itching
- yellowing of the white parts of your eyes, your nails or your skin
- passing dark urine
- passing stools that are very pale

These are signs of a problem with your liver, which can become serious. Your health care provider may carry out blood tests to check for liver problems.

During treatment with [NT005 trade name] you should also tell your health care provider immediately:

- if you develop **severe headache, nausea and vomiting, fits or problems with your vision**.

These may be signs of swelling in your brain or damage to your eyes and you may need to start appropriate treatment straightaway.

Blood tests

If you are having a longer course of treatment with [NT005 trade name] (usually for the tapeworm infection) your health care provider will test your blood before and during your treatment with [NT005 trade name]. This is to check for liver problems and because [NT005 trade name] may reduce the production of blood cell. Your health care provider may decide to stop your treatment if you develop serious liver problems or your blood cell counts decrease too much.

Other medicines and [NT005 trade name]

Tell your health care provider if you are taking, have recently taken or might take any other medicines. This includes herbal medicines and medicines obtained without a prescription.

In particular, tell your health care provider if you are taking:

- Dexamethasone (a corticosteroid)
- Praziquantel (for worm infections)
- Cimetidine (for heartburn)
- Ritonavir (for HIV infection)
- Carbamazepine, phenobarbital, phenytoin, (for epilepsy)

These medicines may affect the action of [NT005 trade name] and your health care provider may need to adjust your treatment.

Pregnancy and contraception

Tell your health care provider if you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby. Ask your health care provider for advice about using effective contraception during a course of treatment; you may need to use contraception for up to 1 month after the end of treatment.

There is not enough information to be certain whether [NT005 trade name] might harm an unborn child, especially with a long course of treatment or if you take it in the first 3 months of pregnancy. If you are pregnant or think you might be, ask your health care provider for advice about taking your medicine. In some cases, your health care provider may suggest delaying treatment with [NT005 trade name] or using another treatment.

It is usually OK to breast feed your baby during treatment with this medicine – ask your health care provider.

Driving and using machines

[NT005 trade name] is not likely to affect your ability to drive or operate machinery.

However, make sure you feel well enough to take on any skilled tasks.

This medicine contains lactose. Lactose is a source of glucose and galactose. The small amount of lactose in each dose may cause symptoms of intolerance, such as diarrhoea, bloated stomach and stomach pains. If, however, you have one of the rare genetic disorders galactosaemia, glucose-galactose intolerance or congenital lactase deficiency you must talk to your health care provider before taking this medicine.

This medicine may contain traces of cow's milk proteins. If you are allergic to cow's milk, talk to your health care provider before taking this medicine.

[NT005 trade name] contains FD&C Yellow #6/Sunset Yellow FCF Aluminium Lake. This may cause allergic reactions.

3. How to take [NT005 trade name]

Always take this medicine exactly as your health care provider has told you. Check with your health care provider if you are not sure.

[NT005 trade name] is available as tablets to be taken by mouth.

For young children or patients not able to swallow the tablets whole, the tablets may be crushed and added to a small amount of semi-solid food or liquid, all of which should be swallowed immediately.

Your health care provider will tell you how many tablets you need to take and for how long, depending on the infection you need to treat. The health care provider will also tell you whether to take your dose before food or with meals.

Use in adults

For the tapeworm infections *cystic echinococcosis* and *alveolar echinococcosis*, the usual dose of [NT005 trade name] is 1 tablet twice daily.

Your health care provider will tell you how long you need to continue treatment, which may be for months or years, depending on where the infection is and what other treatments you are given. Your daily dose may be lower if you weigh less than 60 kg.

For *neurocysticercosis* the usual dose of [NT005 trade name] is 1 tablet twice daily for 10 to 14 days but your health care provider may recommend you take the medicine for longer.

In community programmes to control tapeworms in the gut (*taeniasis*) [NT005 trade name] is given as 1 tablet daily for 3 days.

When [NT005 trade name] is used as part of a community programme to control *lymphatic filariasis* or roundworms in the gut (*ascariasis*, *trichuriasis*, *hookworm*), it is taken as a single tablet once or twice a year.

Use in children

There is limited information on the use of [NT005 trade name] for tapeworm infections in children. In community programmes to control tapeworms in the gut (*taeniasis*) children from 2 years of age may be given 1 tablet of [NT005 trade name] daily for 3 days

For roundworm infections, [NT005 trade name] can be used from the age of 1 year. In children aged 1 to 2 years, the usual dose is half a tablet once or twice a year. In children aged above 2 years, the dose is the same as for adults.

If you take more [NT005 trade name] than you should

If you take too much of the medicine, contact your health care provider straight away for advice. Take your medicine pack and this leaflet with you if you can.

If you forget to take [NT005 trade name]

If you forget to take your dose, take it as soon as you remember.

If you have any further questions on the use of this medicine, ask your health care provider.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

[NT005 trade name] may affect your liver and blood cells. If you are treated with [NT005 trade name] for several days or longer, you will have tests for this before and during treatment.

Speak with your health care provider straightaway if you are taking a course of treatment with this medicine and you get any signs of **liver problems** (see Section 2, Warnings and precautions, for how to recognise these).

Side effects that can occur when treated with [NT005 trade name] for a short time

Common side effects: may affect up to 1 in 10 people

- belly pain, feeling or being sick

Uncommon side effects: may affect up to 1 in 100 people

- headache, dizziness
- diarrhoea
- itchiness, skin rashes

Rare side effects: may affect up to 1 in 1000 people

- allergic reactions (with signs including rash and itchiness)
- bone pain
- protein in the urine
- blood test showing low red cell count
- blood tests showing raised liver enzymes (sign of stress on the liver)

Very rare side effects: may affect up to 1 in 10 000 people

- serious blistering skin rashes with swelling and peeling (erythema multiforme, Stevens-Johnsons syndrome)

Side effects that can occur when treated with [NT005 trade name] for a longer time

Very common: may affect more than 1 in 10 people

- headache
- blood tests showing raised liver enzymes (sign of stress on the liver)

Common side effects: may affect up to 1 in 10 people

- dizziness
- belly pain, feeling or being sick
- temporary thinning of hair and hair loss
- fever

Uncommon side effects: may affect up to 1 in 100 people

- headache, dizziness
- diarrhoea
- itchiness, skin rashes
- hepatitis (inflammation of the liver)
- blood test showing low white cell count
- allergic reactions (with signs including rash, itchiness and urticaria)

Rare side effects: may affect up to 1 in 1000 people

- bone pain
- protein in the urine
- blood test showing low red cell count

Very rare side effects: may affect up to 1 in 10 000 people

- serious blistering skin rashes with swelling and peeling (erythema multiforme, Stevens-Johnsons syndrome)
- blood test showing low count of all blood cells

If any side effects get serious or if you notice any side effects not listed in this leaflet, please tell your health care provider.

Reporting of side effects

If you get a side effect, talk to your health care provider. This includes side effects not listed in this leaflet. You may also be able to report such effects directly to your national reporting system if one is available. By reporting side effects, you can help to improve the available information on this medicine.

5. How to store [NT005 trade name]

Do not store above 30°C. Discard the product 30 days after initial opening

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date stated on the label after EXP. The expiry date refers to the last day of that month.

Do not use this medicine if you notice visible signs of deterioration that it is different from the description below.

Do not throw away any medicines in wastewater or household waste. Ask your health care provider how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What [NT005 trade name] contains

- The active ingredient is albendazole
- The other ingredient(s) of [NT005 trade name] are: lactose monohydrate, corn starch, sodium starch glycolate, sodium lauryl sulphate, povidone, microcrystalline cellulose, saccharin sodium, magnesium stearate and colour FD & C yellow #6 (lake sunset yellow FCF)

There is too little sodium in this medicine to have any effect, even if you are on a low-sodium diet.>

What [NT005 trade name] looks like and contents of the pack

[NT005 trade name] are mottled, pale orange, oblong-shaped uncoated tablets. They are biconvex (rounded on top and bottom), with a break line on one side and “C541” on the other side.

[NT005 trade name] are packed in a white plastic (HDPE) bottle containing 60 tablets. The bottle has a blue child-resistant polypropylene cap and, a heat seal liner and rayon sanyi coil.

Supplier and Manufacturer

Supplier

Cipla Limited
Cipla House
Peninsula Business park
Ganpatrao Kadam Marg
Lower Parel
Mumbai 400 013
India
Tel.: 91 22 4191 6000
Fax : 91 22 4191 6120
Email: globalra@cipla.com

Manufacturer

Cipla Limited Indore (Unit-IV)
Plot No 9, 10
Indore Special Economic Zone, Phase II
Pithampur, Dhar District
Madhya Pradesh 454 775
India

For any information about this medicine, contact the supplier:

This leaflet was last revised in February 2026

Detailed information on this medicine is available on the World Health Organization (WHO) website:
<https://extranet.who.int/prequal/medicines/prequalified/finished-pharmaceutical-products>