

## WHO-PQ RECOMMENDED SUMMARY OF PRODUCT CHARACTERISTICS

*This summary of product characteristics focuses on uses of the medicine covered by WHO's Prequalification Team - Medicines. The recommendations for use are based on WHO guidelines and on information from stringent regulatory authorities.\**

*The medicine may be authorised for additional or different uses by national medicines regulatory authorities.*

---

\*[https://extranet.who.int/prequal/sites/default/files/document\\_files/75%20SRA%20clarification\\_Feb2017\\_newtempl.pdf](https://extranet.who.int/prequal/sites/default/files/document_files/75%20SRA%20clarification_Feb2017_newtempl.pdf)

## 1. NAME OF THE MEDICINAL PRODUCT

[NT016 trade name]<sup>†</sup>

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each chewable tablet contains 400 mg of albendazole.

*Excipients with potential clinical effect* (See section 4-4).

Each chewable tablet also contains:

210 mg of lactose monohydrate

1 mg lake of sunset yellow (FD&C yellow #6)

0.005 mg benzyl alcohol and 0.02 mg sodium benzoate (both from the flavourants)

For the full list of excipients, see section 6-1.

## 3. PHARMACEUTICAL FORM

Chewable tablets.

Light orange, oval, uncoated, mottled tablets. They are biconvex (rounded on top and bottom) with a flat edge. The tablets have a break line on one side and are plain on the other side.

The tablet can be divided into two equal doses.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

[NT016 trade name] is a broad-spectrum anthelmintic used for the treatment of the following infections:

#### **Cestode infections (tapeworms)**

[NT016 trade name] is indicated for the treatment of *Echinococcus multilocularis* and *E. granulosus* infections before or after surgery or where surgery is not suitable.

[NT016 trade name] is further indicated for the treatment of neurocysticercosis caused by larval forms of the pork tapeworm, *Taenia solium*. It may also be given for preventive chemotherapy of *Taenia solium* taeniasis in endemic populations, where other alternatives are not available.

#### **Lymphatic filariasis**

[NT016 trade name] is indicated together with ivermectin and/or diethylcarbamazine for the elimination of lymphatic filariasis.

Treatment is given to the entire eligible population in endemic areas through a mass drug administration programme.

---

<sup>†</sup> Trade names are not prequalified by WHO. This is the national medicines regulatory agency's responsibility.

## Other nematode infections (roundworms)

Albendazole is effective for the treatment of various nematode infections. [NT016 trade name] can be used, alone or in combination with other medicines, for the control of soil-transmitted helminthiasis (ascariasis, trichuriasis and hookworm infections) through mass drug administration programmes.

### 4.2 Posology and method of administration

#### *Posology*

#### **Cestode infections (tapeworms)**

##### *Adults*

In adults over 60 kg, the usual dose for treatment of echinococcosis or *Taenia solium* neurocysticercosis is 400 mg twice a day. In adults up to 60 kg, the dose is 15 mg/kg daily in 2 divided doses (maximum 800 mg daily).

For *cystic echinococcosis*, [NT016 trade name] treatment is continued for 3 to 6 months or longer.

For *alveolar echinococcosis* treatment should be given for at least 2 years and may be continued for many years, reviewed at 2-year intervals.

For *neurocysticercosis*, treatment with [NT016 trade name] is usually for 10–14 days but can be increased to up to 30 days or more for extraparenchymal cysts (e.g. in the ventricles or subarachnoid space).

##### *Children*

Only limited data are available on the use of [NT016 trade name] in children for cestode infections.

##### *Mass drug administration*

In mass drug administration programmes for preventive chemotherapy of *taeniasis* where other alternatives are not available, [NT016 trade name] may be given to endemic populations from 2 years of age in a dose of 400 mg daily for 3 consecutive days. Because of the risk of triggering latent neurocysticercosis, a reporting system must be in place with active surveillance and referral of any neurological adverse events.

#### **Nematode infections**

For the elimination of *lymphatic filariasis* and the control of *soil-transmitted helminthiasis* (ascariasis, trichuriasis, or hookworm disease), a single oral dose of [NT016 trade name] is normally given once a year in mass treatment programmes. The dose may be given twice a year at 6-monthly intervals, if required, in line with national treatment plans. For lymphatic filariasis, [NT016 trade name] is given together with diethylcarbamazine, or ivermectin, or both.

##### *Adults and children aged over 2 years*

In adults and children aged over 2 years, the dose of [NT016 trade name] for mass drug administration is 400 mg.

##### *Children aged 1–2 years*

In children aged 1–2 years, the dose of [NT016 trade name] for the control of soil-transmitted helminthiasis is 200 mg (half a tablet).

[NT016 trade name] is not used in children below 2 years for the elimination of lymphatic filariasis.

#### **Special populations**

##### *Renal impairment*

No dose adjustment is required.

##### *Hepatic impairment*

Caution should be used if [NT016 trade name] is given to patients with liver disease, since albendazole is metabolised by the liver and has been associated with idiosyncratic hepatotoxicity.

### ***Method of administration***

Oral use.

The tablet can be chewed before swallowing.

[NT016 trade name] should be taken with a meal for the prevention and treatment of tissue infections such as echinococcosis, neurocysticercosis and lymphatic filariasis. Taking albendazole with a fatty meal improves its bioavailability and leads to higher blood levels (see section 5.2).

For treatment of intestinal infections such as taeniasis or soil-based helminth infections, [NT016 trade name] should be taken at least 2 hours after a meal and 30 minutes before the next meal. This leads to higher intestinal and lower systemic albendazole levels, which is desirable in these conditions.

### **4.3 Contraindications**

Hypersensitivity to the active substance, other benzimidazoles or to any excipients listed in section 6.1.

### **4.4 Special warnings and precautions for use**

#### *Pre-existing neurocysticercosis*

Treatment with albendazole may uncover pre-existing neurocysticercosis, particularly in areas where taeniasis is common.

Patients may experience neurological symptoms such as seizures, increased intracranial pressure and focal signs as a result of an inflammatory reaction caused by death of the parasite within the brain. Symptoms may develop rapidly after treatment, and appropriate corticosteroid and anticonvulsant therapy should be given straight away.

#### *Risk of retinal damage in patients with retinal neurocysticercosis*

Cysticercosis may, in rare cases, involve the retina. Before starting therapy for neurocysticercosis, the patient should be examined for the presence of retinal lesions. If such lesions are present, the need for anticycsticercal therapy should be weighed against the possibility of retinal damage caused by albendazole-induced changes to the retinal lesion.

#### *Hepatic effects*

Patients undergoing treatment for echinococcosis should have their liver function tested before the start of treatment and regularly (ideally every 2 weeks) during treatment. Patients with disturbed liver function tests before starting albendazole should be carefully evaluated, since the medicine is metabolised by the liver and has been associated with idiosyncratic hepatotoxicity.

Mild to moderate elevations of liver enzymes have been reported frequently with prolonged albendazole treatment. Enzyme abnormalities are usually reversible on discontinuation of treatment. In prolonged higher-dose albendazole therapy for echinococcosis there have been rare reports of severe hepatic abnormalities such as jaundice and histological hepatocellular damage, which may be irreversible.

If enzymes are significantly increased (greater than twice the upper limit of normal) during treatment, [NT016 trade name] should be discontinued. [NT016 trade name] treatment may be reinstituted when levels have returned to normal limits, but liver function should be monitored frequently during repeat therapy.

#### *Bone marrow suppression*

Albendazole can cause bone marrow suppression and therefore blood counts are needed at the start and ideally every two weeks thereafter during treatment for echinococcosis. Patients with liver disease, including hepatic echinococcosis, may be more susceptible to bone marrow suppression leading to pancytopenia, aplastic anaemia, agranulocytosis and leucopenia and therefore warrant closer monitoring of blood counts. Albendazole should be discontinued if clinically significant decreases in blood cell counts occur.

### Excipients

#### *Benzyl alcohol*

Medicines containing benzyl alcohol should not be given to newborn babies (up to 4 weeks old), unless recommended by a doctor. See section 4.2 for information on which children can be given [NT016 trade name]. Benzyl alcohol has been linked with the risk of severe side effects including breathing problems (called “gasping syndrome”) in young children.

[NT016 trade name] should not be used for more than a week in young children (less than 3 years old), unless advised by the health care provider.

[NT016 trade name] should be used with caution, under the health care provider’s advice, in pregnant women, breast-feeding women, and patients with liver or kidney disease. This is because large amounts of benzyl alcohol can build-up in the body, resulting in metabolic acidosis.

Benzyl alcohol may cause allergic reactions.

#### *Sodium benzoate*

Medicines containing sodium benzoate should not be given to newborn babies (up to 4 weeks old), unless recommended by a doctor. See section 4.2 for information on which children can be given [NT016 trade name]. Sodium benzoate can increase bilirubin displacement from albumin, resulting in neonatal jaundice which may develop into kernicterus (non-conjugated bilirubin deposits in the brain tissue).

#### *Lactose*

Patients with congenital lactase deficiency, galactosaemia or glucose-galactose intolerance must not be given this medicine unless strictly necessary.

The small amount of lactose in each dose is unlikely to cause symptoms of lactose intolerance in other patients.

#### *Lake of sunset yellow*

Lake of sunset yellow (FD&C yellow #6) may cause allergic reactions.

It is important to consider the contribution of excipients from all the medicines that the patient is taking.

## **4.5 Interaction with other medicinal products and other forms of interaction**

Cimetidine, dexamethasone and praziquantel may all increase the plasma concentration of the active metabolite of albendazole, albendazole sulfoxide.

Carbamazepine, phenobarbital, phenytoin and ritonavir may reduce plasma concentrations of the active metabolite of albendazole. The clinical relevance of this is unknown, but may result in decreased efficacy, especially in the treatment of systemic helminth infections. Patients should be monitored for efficacy and may require alternative dose regimens or therapies.

## **4.6 Fertility, pregnancy and breastfeeding**

### *Women of childbearing potential*

Pregnancy should be avoided in women treated with albendazole. Adequate contraceptive measures should be taken, particularly with prolonged treatment.

### *Pregnancy*

There are no adequate and well-controlled studies of [NT016 trade name] administration in pregnant women. Limited data from inadvertent single-dose administration of albendazole during the first trimester have not demonstrated an increased incidence of congenital anomalies. However, animal studies with albendazole have revealed evidence of teratogenicity in rats and rabbits (see section 5.3), and therefore preventive chemotherapy with [NT016 trade name] is not recommended in the first trimester of pregnancy.

In general, [NT016 trade name] should be used in pregnant women only if there are no alternatives and the potential benefit justifies the potential risk to the fetus.

#### *Breast-feeding*

Albendazole and its active metabolite pass into breast milk in very small amounts; it is generally considered compatible with breast-feeding, particularly in single doses.

#### *Fertility*

There are no data on the effects of [NT016 trade name] on human male or female fertility.

Animal studies indicate no effects of albendazole on fertility (see section 5.3).

### **4.7 Effects on ability to drive and use machines**

[NT016 trade name] is unlikely to affect the ability to drive or operate machinery.

However, patients should be advised to consider if their clinical status, including any undesirable effects of the medicine, allows them to perform skilled tasks safely.

### **4.8 Undesirable effects**

Data from clinical trials and post-marketing surveillance were used to estimate the frequency of adverse events linked to albendazole.

The adverse reactions considered related to albendazole are listed below by body system, organ class and absolute frequency. Frequencies are defined as very common ( $\geq 1/10$ ), common ( $1/100$  to  $1/10$ ), uncommon ( $1/1000$  to  $1/100$ ), rare ( $1/10\,000$  to  $1/1000$ ), and very rare ( $< 1/10\,000$ ).

#### **Short duration of treatment**

##### *Blood and lymphatic system disorders*

Rare	Low red cell count
------	--------------------

##### *Immune system disorders*

Rare	Hypersensitivity reactions including rash, pruritus and urticaria
------	---

##### *Nervous system disorders*

Uncommon	Headache, dizziness
----------	---------------------

##### *Gastrointestinal disorders*

Common	Upper gastrointestinal symptoms (e.g. epigastric or abdominal pain, nausea, vomiting)
Uncommon	Diarrhoea

##### *Hepatobiliary disorders*

Rare	Elevations of hepatic enzymes
------	-------------------------------

##### *Skin and subcutaneous tissue disorders*

Uncommon	Itchiness, skin rashes
Very rare	Erythema multiforme, Stevens-Johnson syndrome

##### *Musculoskeletal and connective tissue disorders*

Rare	Bone pain
------	-----------

##### *Renal and urinary disorders*

Rare	Proteinuria
------	-------------

#### **Longer duration of treatment**

##### *Blood and lymphatic system disorders*

Uncommon	Leucopenia
Rare	Low red cell count
Very rare	Pancytopenia, aplastic anaemia, agranulocytosis

##### *Immune system disorders*

Uncommon	Hypersensitivity reactions including rash, pruritus and urticaria
----------	---

##### *Nervous system disorders*

Very common	Headache
Common	dizziness

*Gastrointestinal disorders*

Common Gastrointestinal disturbances (abdominal pain, nausea, vomiting)

*Hepatobiliary disorders*

Very common Mild to moderate elevations of hepatic enzymes

Uncommon Hepatitis<sup>1</sup>

*Skin and subcutaneous tissue disorders*

Common Reversible alopecia (thinning of hair, and moderate hair loss)

Very rare Erythema multiforme, Stevens-Johnson syndrome

*Musculoskeletal and connective tissue disorders*

Rare Bone pain

*Renal and urinary disorders*

Rare Proteinuria

*General disorders*

Common Fever

<sup>1</sup> With prolonged albendazole treatment for echinococcosis there have also been reports of severe hepatic abnormalities, including jaundice and hepatocellular damage which may be irreversible

***Reporting of suspected adverse reactions***

Health care providers are asked to report adverse reactions that may be linked to a medicine, to the marketing authorisation holder, or, if available, to the national reporting system. Reports of suspected adverse reactions to a medicine are important for the monitoring of the medicine's benefits and risks.

**4.9 Overdose**

In case of overdosage, symptomatic therapy and general supportive measures are recommended.

**5. PHARMACOLOGICAL PROPERTIES**

**5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Anthelmintics, benzimidazole derivatives, ATC code: P02CA03.

***Mechanism of action***

Albendazole is a benzimidazole derivative that causes degenerative alterations in the tegument and intestinal cells of the parasite and blocks their energy production, ultimately leading to immobilisation and death of the parasite. It works by binding to the colchicine-sensitive site of tubulin, thus inhibiting its assembly into microtubules. As cytoplasmic microtubules are critical in promoting glucose uptake, the glycogen stores of the parasites are depleted.

Albendazole exhibits larvicidal, ovicidal and vermicial activity against helminth parasites. At lower doses the anthelmintic action of albendazole is thought to be mainly intra-intestinal. However, at higher doses, sufficient is absorbed and metabolised to the active sulfoxide metabolite to have a therapeutic effect against tissue parasites.

A number of studies have suggested that therapeutic doses of benzimidazoles are only parasitostatic against *E. multilocularis*. Nonetheless, after several years of albendazole treatment, treatment interruption may be considered, in the absence of progression of the lesions assessed by conventional imaging, and indirect assessment of viability using PET/CT. Although it does not provide direct evidence of *E. multilocularis* viability, and recurrence may occur, this technique, together with the follow-up of specific serum antibodies, may support decision-making and follow-up after albendazole withdrawal in highly selected patients.

**5.2 Pharmacokinetic properties**

The absorption characteristics of [NT016 trade name] have been determined as follows, after administration of one tablet of albendazole 400 mg chewable tablet in healthy volunteers in the fed state:

Pharmacokinetic variable	Albendazole Mean value* ( $\pm$ standard deviation)
Maximum concentration ( $C_{\max}$ )	103 $\pm$ 92 ng/mL
Area under the curve ( $AUC_{0-\infty}$ ), a measure of the extent of absorption	393 $\pm$ 405 ng·h/mL
Time to attain maximum concentration ( $T_{\max}$ ) <sup>#</sup>	2.67 $\pm$ (1.33 – 5.00) h

\*arithmetic mean, <sup>#</sup>median (range)

### Pharmacokinetics of albendazole

<b>General</b>	
	Albendazole concentrations are negligible or undetectable in plasma as it is rapidly converted into the sulfoxide metabolite prior to reaching the systemic circulation. The systemic anthelmintic activity has been attributed to this primary metabolite, albendazole sulfoxide.  Maximal plasma concentrations of albendazole sulfoxide are typically achieved 2 to 5 hours after dosing.
<b>Absorption</b>	
Absolute bioavailability	Not available
Oral bioavailability	Albendazole is poorly absorbed from the gastrointestinal tract (< 5%) due to its low aqueous solubility.
Food effect	Absorption is significantly enhanced (approximately 5-fold) if albendazole is taken with a fatty meal. Following a single 400-mg oral dose of albendazole, the maximum plasma concentration of albendazole sulfoxide was 0.4–1.6 $\mu$ mol/L in fasting patients and 1.8–6.0 $\mu$ mol/L when taken with breakfast (estimated fat content 40 g).
<b>Distribution</b>	
Volume of distribution (mean)	Not available
Plasma protein binding <i>in vitro</i>	Albendazole sulfoxide is 70% bound to plasma protein.
Tissue distribution	Albendazole sulfoxide is widely distributed throughout the body; it has been detected in urine, bile, liver, cyst wall, cyst fluid, and cerebral spinal fluid.
<b>Metabolism</b>	
	Albendazole rapidly undergoes extensive first-pass metabolism in the liver to albendazole sulfoxide, and is generally not detected in plasma. Albendazole sulfoxide is further metabolised to albendazole sulfone and other primary oxidative metabolites.
Active metabolite(s)	Albendazole sulfoxide
<b>Elimination</b>	



Elimination half life	The terminal elimination half-life of albendazole sulfoxide typically ranges from 8 to 12 hours.
Mean systemic clearance (Cl/F)	
Excretion	Following oral administration, albendazole has not been detected in human urine. Albendazole sulfoxide and its metabolites appear to be principally eliminated in bile, with only a small proportion appearing in the urine.
<b>Pharmacokinetic linearity</b>	
	Plasma concentrations of albendazole sulfoxide increase in a dose-proportional manner over the therapeutic dose range following ingestion of a fatty meal.
<b>Drug interactions (<i>in vitro</i>)</b>	
Transporters	Not available
Metabolising enzymes	Not available

## Special populations

### *Renal impairment*

Since renal elimination of albendazole and its primary metabolite, albendazole sulfoxide, is negligible, it is unlikely that clearance of these compounds would be altered in these patients.

### *Liver impairment*

In patients with evidence of extrahepatic obstruction, the systemic availability of albendazole sulfoxide was increased 7-fold.

### *Elderly patients*

Although no studies have investigated the effect of age on albendazole sulfoxide pharmacokinetics, data suggest that the pharmacokinetics are similar to those in young healthy subjects.

### *Children*

Following single-dose administration of 200 to 300 mg (approximately 10 mg/kg) albendazole to paediatric patients with hydatid cyst disease (age range 6 to 13 years), albendazole sulfoxide pharmacokinetics were similar to those observed in fed adults.

## 5.3 Preclinical safety data

### *General toxicity*

Studies of up to 6 months in mice, rats and dogs recognised the haematopoietic system and the liver as target organs of toxicity.

### *Genotoxicity*

In genotoxicity tests, albendazole was found negative in an Ames Salmonella/microsome plate mutation assay, Chinese hamster ovary chromosomal aberration test, and *in vivo* mouse micronucleus test. In the *in vitro* BALB/3T3 cells transformation assay, albendazole produced weak activity in the presence of metabolic activation while no activity was found in the absence of metabolic activation.

### *Carcinogenicity*

Long-term carcinogenicity studies in mice and rats found no evidence of increased incidence of tumours was found in the mice or rats at up to 400 mg/kg/day and 20 mg/kg/day, respectively.

### *Effects on reproduction*

Albendazole did not affect male or female fertility in the rat at an oral dose level of 30 mg/kg/day.

Albendazole was teratogenic (embryotoxicity and skeletal malformations) in pregnant rats and rabbits. The teratogenic response in the rat occurred at oral doses of 10 and 30 mg/kg/day during gestation days 6 to 15, and in pregnant rabbits at oral doses of 30 mg/kg/day during gestation days 7 to 19. In the rabbit study, maternal toxicity (33% mortality) occurred at 30 mg/kg/day. In mice, no teratogenic effects were observed at oral doses up to 30 mg/kg/day administered during gestation days 6 to 15.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Lactose monohydrate  
Microcrystalline cellulose  
Maize starch  
Croscarmellose sodium  
Povidone  
Saccharin sodium  
Sodium lauryl sulfate  
Magnesium stearate  
Colour lake of sunset yellow / FD&C yellow #6  
Flavour trusil orange  
Flavour trusil vanilla

This medicine is essentially 'sodium-free'. It contains less than 1 mmol sodium (23 mg) per tablet.

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

36 months

### **6.4 Special precautions for storage**

Do not store above 30°C. Avoid excursions above 30°C. Protect from light.

### **6.5 Nature and contents of container**

*PVC/PVDC-Alu Blister:*

Clear colourless plastic (PVC/PVDC) on aluminium foil blister cards, each containing 14 tablets. Available in cartons of 4 x 14 tablets.

*Alu/Alu Blister:*

Aluminium foil blister cards, each containing 10 tablets. Available in cartons of 6 x 10 tablets.

### **6.6 Special precautions for disposal and other handling**

No special precautions.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## **7. SUPPLIER**

Indoco Remedies Limited  
166, C.S.T. Road  
Mumbai 400 098, India

Tel. No. +91 22 2087 3310, +91 22 6278 4200

Email: [asenpal@indoco.com](mailto:asenpal@indoco.com)

## 8. WHO REFERENCE NUMBER (WHO Prequalification Programme)

NT016

## 9. DATE OF PREQUALIFICATION

30 August 2024

## 10. DATE OF REVISION OF THE TEXT

January 2025

Section 6 was updated in July 2025

### References

Guideline: alternative mass drug administration regimens to eliminate lymphatic filariasis. Geneva: World Health Organization; 2017 (<https://apps.who.int/iris/bitstream/handle/10665/259381/9789241550161-eng.pdf;jsessionid=4005F8620887C5F63A943EC55982B648?sequence=1>, accessed 15 Sept 2024).

Guideline: preventive chemotherapy to control soil-transmitted helminth infections in at-risk population groups. Geneva: World Health Organization; 2017 (<https://www.who.int/publications/i/item/9789241550116>, accessed 15 Sept 2024)

Guideline for preventive chemotherapy for the control of *Taenia solium* taeniasis. Washington DC: Pan-American Health Organization; 2021 ([https://iris.paho.org/bitstream/handle/10665.2/54800/9789275123720\\_eng.pdf](https://iris.paho.org/bitstream/handle/10665.2/54800/9789275123720_eng.pdf), accessed 15 Sept 2024)

WHO guidelines on management of *Taenia solium* neurocysticercosis. Geneva: World Health Organization; 2021 (<https://iris.who.int/bitstream/handle/10665/344802/9789240032231-eng.pdf>, accessed 15 Sept 2024)

WHO. Considerations for the use of anthelmintic therapy for the treatment of neurocysticercosis. Geneva: World Health Organization; 2023 (<https://iris.who.int/bitstream/handle/10665/366647/9789240068155-eng.pdf>, accessed 26 October 2024)

WHO Fact sheet: Echinococcosis. Geneva: World Health Organization; 2021 (<https://www.who.int/news-room/fact-sheets/detail/echinococcosis>, accessed 15 Sept 2024)

Brunetti E, Kern P, Vuitton DA; Writing Panel for the WHO-IWGE. Expert consensus for the diagnosis and treatment of cystic and alveolar echinococcosis in humans. *Acta Tropica* 2010; 114: 1-16.

<https://doi.org/10.1016/j.actatropica.2009.11.001>, accessed 20 Sept 2024. Zentel 200 mg chewable tablets: Australian product information. Therapeutic Goods Administration; 05/11/2021

<https://www.ebs.tga.gov.au/ebs/picmi/picmirepository.nsf/pdf?OpenAgent&id=CP-2013-PI-01049-1&d=202101161016933>, accessed 15 Sept 2024)

Eskazole 400 mg chewable tablets: Australian product information. Therapeutic Goods Administration; 11/06/2024 (<https://www.ebs.tga.gov.au/ebs/picmi/picmirepository.nsf/pdf?OpenAgent&id=CP-2013-PI-01018-1&d=202011211016933>, accessed September 2024)

Zentel 400 mg compresse: summary of product characteristics. AIFA; 16/10/2021 (<https://medicinali.aifa.gov.it/#/it/organizzazione/882/farmaci/27096/stampati/RCP>, accessed 15 Sept 2024)

### Section 4.6

Albendazole. Drugs and lactation database [online database]. Bethesda: National Institute of Child Health and Human Development; 2024 (<https://www.ncbi.nlm.nih.gov/books/NBK501368/>, accessed 15 September 2024)

Detailed information on this medicine is available on the World Health Organization (WHO) website: <https://extranet.who.int/prequal/medicines/prequalified/finished-pharmaceutical-products>

