

## WHO-PQ RECOMMENDED SUMMARY OF PRODUCT CHARACTERISTICS

*This summary of product characteristics focuses on uses of the medicine covered by WHO's Prequalification Team - Medicines. The recommendations for use are based on WHO guidelines and on information from stringent regulatory authorities.\*  
The medicine may be authorised for additional or different uses by national medicines regulatory authorities.*

---

\*[https://extranet.who.int/pqweb/sites/default/files/documents/75%20SRA%20clarification\\_Feb2017\\_newtempl.pdf](https://extranet.who.int/pqweb/sites/default/files/documents/75%20SRA%20clarification_Feb2017_newtempl.pdf)

## 1. NAME OF THE MEDICINAL PRODUCT

[RH031 trade name]†

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains 1.5 mg levonorgestrel.

*Excipient with known effect:*

Each tablet also contains 154 mg of lactose monohydrate.

For a full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Tablet.

Round, white to off-white, uncoated flat tablet, debossed '145' on one side and plain on the other side.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Emergency contraception within 72 hours of unprotected sexual intercourse or failure of a contraceptive method.

### 4.2 Posology and method of administration

For oral administration. The treatment comprises a single tablet that should be taken as soon as possible.

The highest efficacy is achieved if the tablet is taken as soon as possible, preferably within 12 hours (and no later than 72 hours) after unprotected intercourse. (For efficacy data, see section 5.1.)

[RH031 trade name] can be used at any time during the menstrual cycle unless menstrual bleeding is overdue.

If vomiting occurs within three hours of taking the tablet, another tablet should be taken immediately. If repeated vomiting occurs, the tablet may be administered vaginally.

Women who have used enzyme-inducing drugs during the last 4 weeks and need emergency contraception are recommended to take a double dose of levonorgestrel (i.e. 2 tablets taken together).

After using emergency contraception, it is recommended to use a local barrier method (condom, diaphragm, spermicide, cervical cap) until the next menstrual period starts. The use of [RH031 trade name] does not contraindicate the continuation of regular hormonal contraception.

[RH031 trade name] is not recommended for use by young women aged under 16 years without medical supervision.

### 4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

### 4.4 Special warnings and precautions for use

Emergency contraception is not effective in terminating an existing pregnancy.

Emergency contraception is an occasional method. It should **not** replace a regular contraceptive method.

Efficacy appears to decline with time (see section 5.1).

---

† Trade names are not prequalified by WHO. This is the national medicines regulatory agency's responsibility.

If there is uncertainty about the timing of the unprotected intercourse or if the woman has had unprotected intercourse more than 72 hours earlier in the same menstrual cycle, conception may have occurred. Treatment with [RH031 trade name] following the second act of intercourse may therefore be ineffective in preventing pregnancy. If menstrual periods are delayed by more than 5 days or abnormal bleeding occurs at the expected date of menstrual periods or pregnancy is suspected for any other reason, pregnancy should be ruled out.

If pregnancy occurs after treatment with [RH031 trade name], the possibility of an ectopic pregnancy should be considered, especially in women in whom severe abdominal pain or fainting occurs, or if there is a history of ectopic pregnancy, Fallopian tube surgery or pelvic inflammatory disease. The absolute risk of ectopic pregnancy is likely to be low, as levonorgestrel prevents ovulation and fertilisation. Ectopic pregnancy may continue despite uterine bleeding. Therefore, [RH031 trade name] is not recommended for women at risk of ectopic pregnancy (history of salpingitis or of ectopic pregnancy).

[RH031 trade name] is not recommended in patients with severe hepatic dysfunction.

Severe malabsorption syndromes, such as Crohn's disease, might impair the efficacy of [RH031 trade name].

After taking [RH031 trade name], menstrual periods are usually normal and occur at the expected date. They can sometimes occur earlier or later than expected by a few days. Women should be advised to see a health care provider to initiate or adopt a method of regular contraception. If no withdrawal bleed occurs in the next pill-free period following the use of [RH031 trade name] after regular hormonal contraception, pregnancy should be ruled out.

Repeated administration within a menstrual cycle is not advisable because of the possibility of disturbing the cycle.

Any regular contraceptive method can be started immediately after the use of [RH031 trade name] emergency contraceptive pills. If the woman starts a hormonal contraceptive:

- she needs to abstain from sexual intercourse or use barrier contraception for 7 days
- she should be advised to have a pregnancy test if she does not have a withdrawal bleed within 3 weeks.

Limited and inconclusive data suggest that there may be reduced efficacy of levonorgestrel with increasing body weight or body mass index (BMI) (see sections 5.1 and 5.2). In all women, emergency contraception should be taken as soon as possible after unprotected intercourse, regardless of the woman's body weight or BMI.

[RH031 trade name] is not as effective as a conventional regular method of contraception and is suitable only as an emergency measure. Women who present for repeated courses of emergency contraception should be advised to consider long-term methods of contraception.

Use of emergency contraception does not replace the necessary precautions against sexually transmitted diseases.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

The metabolism of levonorgestrel is enhanced by concomitant use of liver enzyme inducers, mainly CYP3A4 enzyme inducers.

Drugs suspected of having the capacity to reduce the efficacy of levonorgestrel include barbiturates (including primidone), phenytoin, carbamazepine, herbal medicines containing St. John's wort (*Hypericum perforatum*), rifampicin, ritonavir, rifabutin, bosentan, felbamate, oxcarbazepine and griseofulvin.

Significant changes (increase or decrease) in the plasma levels of the progestogen have been noted in some cases of co-administration with HIV protease inhibitors or with non-nucleoside reverse transcriptase inhibitors. The potential interaction may require close monitoring, alteration of drug dosage or timing of administration.

For women who have used enzyme-inducing drugs in the past 4 weeks and need emergency contraception, a double-dose of levonorgestrel should be taken (see section 4.2).

<sup>#</sup> Formerly Famy Care Ltd then Jai Pharma Ltd

Medicines containing levonorgestrel may increase the risk of cyclosporin toxicity due to possible inhibition of cyclosporin metabolism.

#### 4.6 Fertility, pregnancy and breastfeeding

##### *Pregnancy*

[RH031 trade name] should not be given to pregnant women. It will not interrupt the pregnancy.

In case of failure of this emergency contraception and developing pregnancy, epidemiological studies indicate no adverse effects of progestogens on the fetus. There are no clinical data on the potential consequences if doses greater than 1.5 mg levonorgestrel are taken (see section 5.3).

##### *Breast-feeding*

Levonorgestrel is secreted into breast milk. Potential exposure of an infant to levonorgestrel can be reduced if the breast-feeding woman takes the tablets immediately after feeding and avoids nursing following each [RH031 trade name] administration.

##### *Fertility*

Levonorgestrel increases the possibility of menstrual cycle disturbances which can sometimes lead to earlier or later ovulation date. These changes can result in modified fertility date; however there are no fertility data in the long term.

#### 4.7 Effects on ability to drive and use machines

No studies on the effect on the ability to drive and use machines have been performed.

#### 4.8 Undesirable effects

The most commonly reported undesirable effect was nausea.

All adverse drug reactions are listed by system, organ class and frequency.

Frequencies are defined as very common ( $\geq 1/10$ ), common ( $\geq 1/100, <1/10$ ), uncommon ( $\geq 1/1,000, <1/100$ ), rare ( $\geq 1/10,000, <1/1,000$ ), very rare ( $<1/10,000$ ), not known (cannot be estimated from the available data).

System Organ Class	Frequency of adverse reactions		
	Very common	Common	Very rare
<b>Gastrointestinal disorders</b>	Nausea Lower abdominal pain	Diarrhoea Vomiting	
<b>General disorders and administration site conditions</b>	Fatigue		Face oedema
<b>Nervous system disorders</b>	Headache	Dizziness	
<b>Reproductive system and breast disorders</b>	Bleeding not related to menses*	Delay of menses more than 7 days** Menstruation irregular Breast tenderness	Pelvic pain Dysmenorrhoea
<b>Skin and subcutaneous tissue disorders</b>			Rash Urticaria Pruritus

\* Bleeding patterns may be temporarily disturbed, but most women will have their next menstrual period within 7 days of the expected time.

\*\* If the next menstrual period is more than 5 days overdue, pregnancy should be excluded.

### ***Reporting of suspected adverse reactions***

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Health care providers are asked to report any suspected adverse reactions to the marketing authorisation holder, or, if available, via the national reporting system.

### **4.9 Overdose**

There have been no reports of any serious damage to health caused by an overdose.

The symptoms that may occur in such a case include: nausea, vomiting and mild vaginal bleeding. There is no specific antidote. Treatment should be symptomatic.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Sex hormones and modulators of the genital system, emergency contraceptives, ATC code: G03AD01

#### Mechanism of action

The precise mode of action of levonorgestrel is not known.

At the recommended regimen, levonorgestrel is thought to work mainly by preventing ovulation and fertilisation if intercourse has taken place in the pre-ovulatory phase, when the likelihood of fertilisation is the highest. It is not effective once the process of implantation has begun.

#### Clinical efficacy and safety

*Efficacy:* Results from a randomised, double-blind clinical study conducted in 2001 (Lancet 2002; 360: 1803-1810) showed that a 1.5-mg single dose of levonorgestrel (taken within 72 hours of unprotected sex) prevented 84% of expected pregnancies (compared with 79% when two 750-microgram tablets were taken 12 hours apart).

It is therefore, recommended that levonorgestrel is taken as soon as possible (and no later than 72 hours) after unprotected intercourse.

At the recommended regimen, levonorgestrel is not expected to significantly modify blood clotting factors, or lipid and carbohydrate metabolism.

*Safety:* Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

A double-blind, controlled clinical trial in 1,955 evaluable women compared the efficacy and safety of Levonorgestrel (one 0.75 mg tablet of levonorgestrel taken within 72 hours of unprotected intercourse, and one tablet taken 12 hours later) to the Yuzpe regimen (two tablets each containing 250 micrograms levonorgestrel and 50 micrograms ethinylestradiol, taken within 72 hours of intercourse, and two tablets taken 12 hours later).

There are limited and inconclusive data on the effect of high body-weight/BMI on the contraceptive efficacy. In three WHO studies, no trend for a reduced efficacy with increasing body-weight/BMI was observed), whereas in the two other studies (Creinin et al., 2006 and Glasier et al., 2010) a reduced contraceptive efficacy was observed with increasing body weight or BMI. Both meta-analyses excluded intake later than 72 hours after unprotected intercourse (i.e. off-label use of levonorgestrel) and women who had further acts of unprotected intercourse.

#### Paediatric population

A prospective observational study showed that out of 305 treatments with levonorgestrel emergency contraceptive tablets, seven women became pregnant resulting in an overall failure rate of 2.3%. The failure

rate in women under 18 years (2.6% or 4/153) was comparable to the failure rate in women 18 years and over (2.0% or 3/152).

## 5.2 Pharmacokinetic properties

### Absorption of [RH031 trade name]

The absorption characteristics of [RH031 trade name] have been determined after administration of one (1) tablet in healthy volunteers in the fasted state, as follows:

Pharmacokinetic variable	Arithmetic mean $\pm$ standard deviation (*)
	Levonorgestrel
Maximum concentration ( $C_{max}$ )	20.1 $\pm$ 6.6 (19.3) ng/mL
Area under the curve ( $AUC_{0-72h}$ ), a measure of the extent of absorption	318 $\pm$ 138 (289) ng·hour/mL
Time to attain maximum concentration ( $t_{max}$ ) <sup>#</sup>	2.13 (1–4) h

\*geometric mean; # median (range)

### Pharmacokinetics of levonorgestrel

<b>Absorption</b>	
Oral bioavailability	Rapid and near complete absorption
Food effect	NA
<b>Distribution</b>	
Volume of distribution (mean)	106 L
Plasma protein binding <i>in vitro</i>	33.5% bound to serum albumin and 65% bound to sex hormone binding globulin
Tissue distribution	0.1% of dose transferred to breast milk
<b>Metabolism</b>	
General	Metabolism follows the known pathways of steroid metabolism. 100% metabolized by liver.
Active metabolite(s)	None known.
<b>Elimination</b>	
Elimination half life	26 hours
Mean systemic clearance (Cl/F)	1.0 - 1.5 mL/minute/kg
% of dose excreted in urine	Levonorgestrel metabolites are excreted in about equal proportions with urine and faeces.
% of dose excreted in faeces	Levonorgestrel metabolites are excreted in about equal proportions with urine and faeces.

\*Information not available

### Pharmacokinetics in obese women

A pharmacokinetic study showed that levonorgestrel concentrations are decreased in obese women (BMI  $\geq$  30 kg/m<sup>2</sup>) (approximately 50% decrease in  $C_{max}$  and  $AUC_{0-24}$ ), compared to women with normal BMI (< 25 kg/m<sup>2</sup>) (Praditpan et al., 2017). Another study also reported a decrease of levonorgestrel  $C_{max}$  by approximately 50% between obese and normal BMI women, while doubling the dose (3 mg) in obese women appeared to provide plasma concentration levels similar to those observed in normal women who received 1.5 mg of levonorgestrel (Edelman et al., 2016). The clinical relevance of these data is unclear.

### **5.3 Preclinical safety data**

Non-clinical data reveal no special hazard for humans based on conventional studies of chronic toxicity, mutagenicity and carcinogenicity potential, beyond the information included in other sections of the SmPC.

Animal experiments with levonorgestrel have shown virilisation of female foetuses at high doses.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Povidone  
Lactose monohydrate  
Maize starch  
Colloidal silicon dioxide  
Magnesium Stearate

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

36 months

### **6.4 Special precautions for storage**

Do not store above 30°C. Protect from light. Store the tablet in the blister in the provided carton.

### **6.5 Nature and contents of container**

PVC-PVdC/Aluminium blister containing one tablet. One blister card per carton.

### **6.6 Special precautions for disposal and other handling**

Any unused product or waste material should be disposed of in accordance with local requirements.

## **7. SUPPLIER**

Mylan Laboratories Limited  
Plot No. 564/A/22  
Road No. 92 Jubilee Hills  
Hyderabad, Telangana - 500096  
India

Email: [Imtiyaz.basade@viatris.com](mailto:Imtiyaz.basade@viatris.com)

## **8. WHO REFERENCE NUMBER (WHO Prequalification Programme)**

RH031

## **9. DATE OF PREQUALIFICATION**

21 October 2013

## **10. DATE OF REVISION OF THE TEXT**

September 2021

## **References**

### General references

Levonorgestrel SmPC, January 2019 available at

<https://www.medicines.org.uk/emc/product/7308/smpc>

WHO key facts on Emergency Contraception:

<http://www.who.int/en/news-room/fact-sheets/detail/emergency-contraception>

Selected Practice Recommendations for Contraceptive Use, Third Edition 2016, World Health Organization:

[http://www.who.int/reproductivehealth/publications/family\\_planning/SPR-3/en/](http://www.who.int/reproductivehealth/publications/family_planning/SPR-3/en/)

Medical Eligibility Criteria for Contraceptive Use, Fifth Edition 2015. World Health Organization:

[http://www.who.int/reproductivehealth/publications/family\\_planning/Ex-Summ-MEC-5/en/](http://www.who.int/reproductivehealth/publications/family_planning/Ex-Summ-MEC-5/en/)

US Selected Practice Recommendations for Contraceptive Use, 2016

US Centre for Disease Control and Prevention

MMWR , Recommendations and Reports/ July 29, 2016/65(4); 1-66:

<https://www.cdc.gov/mmwr/volumes/65/rr/rr6504a1.htm>

### Sections 4.4 and 5.2

Praditpan et al. Pharmacokinetics of levonorgestrel and ulipristal acetate emergency contraception in women

with normal and obese body mass index. *Contraception* [online] 2017;95(5):464-469. Available from

[https://www.contraceptionjournal.org/article/S0010-7824\(17\)30010-0/fulltext](https://www.contraceptionjournal.org/article/S0010-7824(17)30010-0/fulltext) [Accessed 26 March 2020]

Edelman et al. Impact of obesity on the pharmacokinetics of levonorgestrel-based emergency contraception: single and double dosing. *Contraception* [online] 2016; 94(1):52-7. Available from

<https://dx.doi.org/10.1016%2Fj.contraception.2016.03.006> [Accessed 26 March 2020]

### Section 5.1

Von Hertzen et al. Low dose mifepristone and two regimens of levonorgestrel for emergency contraception:

a WHO multicentre randomised trial. *Lancet* [online] 2002;360(9348):1803-10. Available from

[https://doi.org/10.1016/S0140-6736\(02\)11767-3](https://doi.org/10.1016/S0140-6736(02)11767-3) [Accessed 26 March 2020]

Creinin et al. Progesterone receptor modulator for emergency contraception: a randomized controlled trial.

*Obstet Gynecol* [online] 2006; 108(5), 1089-1097. Available from

10.1097/01.aog.0000239440.02284.45 [Accessed 26 March 2020]

Glasier et al. Ulipristal acetate versus levonorgestrel for emergency contraception: a randomised non-

inferiority trial and meta-analysis. *Lancet* [online] 2010; 13;375(9714):555-62. Available from

10.1016/S0140-6736(10)60101-8 [Accessed 26 March 2020]

*All web links last accessed September 2021.*

*Detailed information on this medicine is available on the World Health Organization (WHO) website:*

<https://extranet.who.int/pqweb/medicines>