

## **WHO-PQ RECOMMENDED PATIENT INFORMATION LEAFLET**

*This patient information leaflet focuses on uses of the medicine covered by WHO's Prequalification Team - Medicines. The recommendations for use are based on WHO guidelines and on information from stringent regulatory authorities.\**  
*The medicine may be authorised for additional or different uses by national medicines regulatory authorities.*

---

\* [https://extranet.who.int/prequal/sites/default/files/document\\_files/75%20SRA%20clarification\\_Feb2017\\_newtempl.pdf](https://extranet.who.int/prequal/sites/default/files/document_files/75%20SRA%20clarification_Feb2017_newtempl.pdf)  
Page 1 of 7

## Information for the patient

[RH083 trade name] <sup>†</sup>  
Oxytocin

*If you are a carer or parent looking after the person who takes this medicine, use this leaflet to give the medicine correctly and take note of the warnings and side effects.*

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have questions about the medicine, ask your health care provider.
- This medicine is for you only. Do not pass it on to others. It may harm them, even if their illness seems to be the same as yours.
- If you get any side effects, talk to your health care provider. This includes unwanted effects not listed in this leaflet. See section 4.

### What is in this leaflet

1. What [RH083 trade name] is and what it is used for
2. What you need to know before you take [RH083 trade name]
3. How to take [RH083 trade name]
4. Possible side effects
5. How to store [RH083 trade name]
6. Contents of the pack and other information

#### 1. What [RH083 trade name] is and what it is used for

[RH083 trade name] contains a manufactured form of oxytocin (a natural hormone). It makes the muscles of the womb contract.

[RH083 trade name] is used:

- To start or help contractions during childbirth (labour)
- To help in the management of miscarriage
- During a caesarian section
- To prevent and control bleeding after delivery of your baby

#### 2. What you need to know before you are given [RH083 trade name]

**You must not receive [RH083 trade name]:**

- If you are allergic to oxytocin or any of the other ingredients of this medicine (listed in section 6)
- If your health care provider thinks that to start or increase contractions of the womb would be unsuitable, for example:
  - if contractions of the womb are unusually strong
  - if there is a blockage that could prevent delivery
  - if your baby may not be receiving enough oxygen
- Where labour or delivery through the birth canal is not advisable, for example, if:
  - your baby's head is too large to fit through the birth canal

---

<sup>†</sup>Trade names are not prequalified by WHO. This is the national medicines regulatory agency's responsibility.

- your baby is not in the right position in the birth canal
- the placenta lies near or over the neck of your womb
- your baby separates from the womb before the baby is born
- there are loops of umbilical cord between the baby and the neck of the womb either before or after your waters break
- your womb is over-stretched and more likely to tear, for example if you are carrying more than one baby or have too much water (amniotic fluid) in your womb
- you have had at least five pregnancies in the past or if your womb is scarred by previous caesarean section or other surgery
- If you have just been given medicines called prostaglandins (used to bring on labour or treat stomach ulcers). [RH083 trade name] should be used at least 6 hours after vaginal prostaglandins as the combined effects of both medicines may be too strong.

[RH083 trade name] should not be used for prolonged periods if:

- the treatment does not increase your contractions
- you have a condition known as severe pre-eclamptic toxemia (high blood pressure, protein in the urine and swelling)
- you have severe problems with your heart or blood circulation.

### **Warnings and precautions**

Talk to your health care provider before you receive [RH083 trade name] if:

- you are prone to chest pain because of problems with your heart or blood circulation
- you have an irregular heartbeat ('long QT syndrome') or problems with your heart rhythm, or if you are taking medicines that affect the heart rhythm (see under 'Other medicines and [RH083 trade name]', below')
- you have previously had a caesarean section
- you are more than 35 years old
- you have raised blood pressure or heart problems
- your womb was contracting strongly but has now begun to contract less strongly
- your health care provider has told you that normal delivery may be difficult because your pelvis is too small
- you have kidney problems, as oxytocin may cause build-up of water in your body
- you have had complications during your pregnancy
- you are more than 40 weeks pregnant

Your health care provider will look out for any complications that can arise from using this medicine.

When this medicine is given into a vein, your health care provider will make sure it is given at the proper speed. Tell your health care provider if you feel faint or dizzy, your heartbeat gets fast, or you feel hot and flushed all over. This can happen if the medicine is given too fast.

When [RH083 trade name] is given to induce and enhance labour, your health care provider will adjust the infusion speed so that you have contractions similar to normal labour. If the speed is too high the contractions may be too strong and too long, which may cause the womb to tear with serious complications for you and your baby.

High doses of [RH083 trade name] may force amniotic fluid from your womb into your blood. This is known as amniotic fluid embolism.

Tell your health care provider if, after your baby is born, you bleed too easily or have anaemia because these may be signs of a rare side effect called disseminated intravascular coagulation.

Large doses of [RH083 trade name] over a long period whilst drinking or receiving large amount of fluid may cause a build-up of water in your body, cause difficulty in breathing and lower salt levels in your blood.

If any of the above apply to you, or if you are not sure, speak to your health care provider before you receive [RH083 trade name].

#### *Latex allergy*

The active substance in [RH083 trade name] can cause a severe allergic reaction (anaphylaxis) in patients with latex (rubber) allergy. Tell your health care provider if you are allergic to latex.

#### *Children and adolescents*

[RH083 trade name] is not intended for use in children or adolescents.

#### *Other medications and [RH083 trade name]*

Tell your health care provider if you are taking or have recently taken any of the following medicines as they may interfere with [RH083 trade name]:

- Prostaglandins (used to start labour or to treat stomach ulcers) and similar medicines as the effects of both drugs may be increased
- Medicines that can cause an irregular heartbeat, as oxytocin may increase this effect
- Anaesthetics which you breathe in (e.g. to put you to sleep during surgery), such as halothane, cyclopropane, sevoflurane or desflurane, as these may weaken your contractions, or cause problems with your heartbeat
- Medicines that can increase your blood pressure or raise your heartbeat (which are sometimes also included in anaesthetic medicines applied to your body to relieve pain). Using these medicines at the same time as [RH083 trade name] may increase blood pressure.

Tell your health care provider if you are taking or have recently taken or might take any other medicines.

#### *[RH083 trade name] with food and drink*

Your health care provider may tell you to keep the amount of fluids you drink to a minimum.

#### *Pregnancy, breastfeeding and fertility*

Based on the wide experience of use and the nature of this medicine, [RH083 trade name] is not expected to harm your baby when used correctly.

Oxytocin may be present in small amounts in breast milk but is not expected to have harmful effects because it is quickly inactivated by your baby's digestive system. The effects of oxytocin on fertility are unknown.

#### *Driving and using machines*

[RH083 trade name] can start labour. Women with uterine contractions should not drive or use machines.

#### **[RH083 trade name] contains ethanol**

This medicine contains small amounts of ethanol (alcohol), less than 100 mg per mL (0.005 mg). The small amount of alcohol in this medicine will not have any noticeable effects.

### **3. How [RH083 trade name] is given**

[RH083 trade name] is given to you by a health care provider in a hospital.

Your health care provider will decide when and how to treat you with [RH083 trade name]. If you think that the effect of [RH083 trade name] is too strong or too weak, tell your health care provider.

While you are receiving [RH083 trade name], you and your baby will be closely monitored.

[RH083 trade name] is usually given as an infusion (drip) into one of your veins but it can also be given by injection into your muscles.

The dose you get depends on what the medicine is being used for.

*To start or help contractions during labour*

The rate of infusion will start at 10 drops per minute. This may be gradually increased to a maximum rate of 60 drops per minute. If necessary, your health care provider may change the infusion bag to increase the amount of medicine in the infusion bag. The infusion rate can often be reduced once you have a good pattern of contractions of about 3 to 4 contractions every 10 minutes.

If you do not reach a good pattern of contractions with the medicine, you may need to have a caesarean section to deliver your baby safely.

*Miscarriage*

The dose is 5 units by infusion into a vein. Your health care provider may continue the infusion if you need more medicine.

*Caesarean section*

The dose is 5 units by infusion into a vein immediately after delivery of your baby.

*Prevention of bleeding after delivery*

The dose is 10 units by injection either into your vein or into a muscle. Your health care provider may give you the medicine into your vein if you are already being given other medicines in this way.

*Treatment of bleeding after delivery*

If you already have a drip running, you may be given 10 to 40 units in the drip. Your health care provider will adjust the speed of the drip depending on your contractions. In some cases, you may be given the medicine by injection into your muscle.

*If you receive more [RH083 trade name] than you should*

As this medicine is given to you by your health care provider, it is very unlikely that you will receive an overdose.

If anyone accidentally receives this medicine, tell the hospital accident and emergency department or a health care provider immediately. Show any left-over medicines or the empty packet to the doctor.

An overdose of [RH083 trade name] could cause:

- Very strong contractions of your womb
- Damage to your womb which could include tearing
- The placenta to come away from your womb
- Amniotic fluid (the fluid around the baby) to enter your bloodstream
- Harm to your baby

If you have any further questions on the use of this medicine, ask your health care provider.

#### **4. Possible side effects**

##### **Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

##### ***Effects in the mother***

*Common side effects* (affects more than 1 in 100 patients):

- Headache
- Fast or slow heartbeat
- Feeling sick (nausea) or being sick (vomiting)

*Uncommon side effects* (affects more than 1 in 1000 patients):

- An irregular heartbeat

*Rare side effects* (affects more than 1 in 10,000 patients):

- Severe allergic reaction with difficulty breathing, dizziness and light-headedness, feeling faint, nausea, cold and clammy skin or a fast or weak pulse
- Rash

*The following side effects can also occur, but it is not known how common they are:*

- Chest pain (angina)
- Irregular heartbeat (QTc prolongation seen on electrocardiogram)
- Low blood pressure
- Haemorrhage (bleeding)
- Excessive contractions
- Excessive or continuous contractions
- Tearing of the womb
- Rapid swelling under the skin in areas such as the face, throat, arms and legs
- Fluid retention (water build-up). Symptoms may include headache, anorexia (loss of appetite), feeling or being sick, stomach pain, sluggishness, drowsiness, unconsciousness, low levels of certain chemicals in the blood (e.g. sodium or potassium), fits
- Low blood salt levels
- Sudden fluid overload in the lungs
- Sudden brief sensation of heat over the whole body
- Abnormal blood clotting, bleeding or anaemia (disseminated intravascular coagulation)

### ***Effects in the baby***

*The following side effects can also occur, but it is not known how common they are:*

- Low blood salt levels
- Excessive contractions in the womb may cause shortage of oxygen, suffocation and death

### **Reporting of side effects**

If you get a side effect, talk to your health care provider. This includes side effects not listed in this leaflet. You may also be able to report such effects directly to your national reporting system if one is available. By reporting side effects, you can help to improve the available information on this medicine.

## **5. How to store [RH083 trade name]**

Keep this medicine out of the sight and reach of children.

Store in a refrigerator (2°C to 8°C) in the original carton, protected from light. Do not freeze.

This medicine must not be used after the expiry date stated on the label after 'EXP'. The expiry date refers to the last day of that month.

This medicine must not be used if there are description of the visible signs of deterioration.

Do not throw away any medicines in wastewater or household waste. Ask your health care provider how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What [RH083 trade name] contains**

- The active ingredient is oxytocin
- The other ingredients of {product name} are ethanol, chlorobutanol, sodium acetate trihydrate, sodium chloride, glacial acetic acid, water for injections

There is too little sodium in this medicine to have any effect, even if you are on a low-sodium diet.

**What [RH083 trade name] looks like and contents of the pack**

[RH083 trade name] is a clear colourless solution, contained in a clear type I glass ampoule.

Pack sizes: 5, 10 ampoules per plastic ampoule tray or 100 ampoules in a honey comb pack.

**Supplier and Manufacturer**

***Supplier***

Steril-Gene Life Sciences (P) Ltd  
No.15, Gopalakrishna Road  
T. Nagar, Chennai-600 017  
India  
Tel: +91 44 23452030 /34  
Fax: +91 44 23452036  
e-mail: vijay.balakumar@steril-gene.com

***Manufacturer***

Steril-Gene Life Sciences (P) Ltd  
45, Mangalam Main Road  
Mangalam Village  
Villianur Commune  
Puducherry- 605110  
India  
Tel: +91 413 2661103  
Fax: +91 413 2661102  
e-mail: saravankumar.v@steril-gene.com

*“For any information about this medicine, contact the supplier”.*

**This leaflet was last revised in March 2025**

Detailed information on this medicine is available on the World Health Organization (WHO) website:  
<https://extranet.who.int/prequal/medicines/prequalified/finished-pharmaceutical-products>