

## WHO-PQ RECOMMENDED SUMMARY OF PRODUCT CHARACTERISTICS

*This summary of product characteristics focuses on uses of the medicine covered by WHO's Prequalification Team - Medicines. The recommendations for use are based on WHO guidelines and on information from stringent regulatory authorities.\**

*The medicine may be authorised for additional or different uses by national medicines regulatory authorities.*

---

\*[https://extranet.who.int/pqweb/sites/default/files/documents/75%20SRA%20clarification\\_Feb2017\\_newtempl.pdf](https://extranet.who.int/pqweb/sites/default/files/documents/75%20SRA%20clarification_Feb2017_newtempl.pdf)

## 1. NAME OF THE MEDICINAL PRODUCT

[TB222 trade name]†

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each hard capsule contains 250 mg cycloserine.

Each capsule shell contains about 57 µg ponceau 4R (cochineal red A). See section 4-4.  
For the full list of excipients, see section 6.1

## 3. PHARMACEUTICAL FORM

Hard Capsules.

Each hard gelatin capsule, with orange-coloured cap and white-coloured body, is filled with white or almost white powder.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

[TB222 trade name] is indicated in combination with other antituberculosis agents for the treatment of drug-resistant tuberculosis due to *Mycobacterium tuberculosis*.

Treatment regimens should follow most recent WHO treatment guidelines, supplemented by other authoritative guidelines.

### 4.2 Posology and method of administration

Oral use

[TB222 trade name] **must always be given in combination** with other antituberculosis agents.

Pyridoxine (vitamin B6) should be taken concomitantly with cycloserine (see section 4.4).

#### Posology

##### Adults:

The usual dose is 10-15 mg/kg/day, up to a usual maximum of 1000 mg/day, given in two divided doses every 12 hours or once a day if tolerated.

Body weight	Daily dose	Number of capsules daily
30-45 kg	500 mg	2
≥46 kg	750 mg	3

##### Children:

The recommended dose is 15–20 mg/kg/day given as a single dose or in two divided doses every 12 hours. A daily dose of 1000 mg should not be exceeded. If available, therapeutic drug monitoring may be useful. Peak concentrations between 15-40 µg/mL have been recommended as appropriate.

Patients under 15 years of age and weighing less than 16 kg, should be given other formulations to allow appropriate dosage, e.g., capsules containing 125 mg of cycloserine.

If such formulations are not available, an extemporaneous preparation may be prepared by dispersing the contents of a 250-mg capsule of [TB222 trade name] in 10 mL of drinking water in order to facilitate administration in patients in lower weight-bands, although bioavailability is uncertain.

For dosing of infants weighing less than 5 kg an expert in treatment of paediatric drug-resistant tuberculosis should be consulted whenever possible.

---

† Trade names are not prequalified by WHO. This is the national medicines regulatory agency's responsibility.

Recommended daily doses are as follows:

Body weight of child	Daily dose	Daily volume* of extemporaneous preparation to be taken
3-4 kg	25 mg	1 mL
5-6 kg	50 mg	2 mL
7-9 kg	125 mg	5 mL
10-15 kg	250 mg	Give as 250-mg capsule**
≥ 16 kg	500 mg	Give as two 250-mg capsules**

**\*If a dose other than 5 or 10 mL is recommended for administration by a caregiver, or the daily dose is to be divided, the health care provider should advise the caregiver on how to measure or divide the dose and ensure a suitable oral syringe is available for measurement**

**\*\*For older children weighing 16 kg or more who cannot swallow capsules, the 250-mg capsules can be opened and dispersed similarly in 10 mL water to aid administration.**

For detailed instructions on preparing an extemporaneous formulation, see below: “Method of administration, extemporaneous formulation for children”

For **tuberculous meningitis** different dosing regimens may apply. Current WHO treatment guidelines should be followed.

#### ***Dose adjustments***

Some patients may require alternate day 250 mg and 500 mg dosing to avoid toxicity.

To minimise headaches at the start of therapy, cycloserine can be started at lower doses of 250–500 mg and gradually increased over one to two weeks to achieve the target dose.

#### ***Renal failure/dialysis:***

For patients with creatinine clearance < 30 mL/min or for patients on haemodialysis the recommended dose is 250 mg once daily or 500 mg, 3 times per week. Doses should be given after haemodialysis. Drug concentrations should be monitored to keep peak concentrations <35 µg/mL. Patients should also be carefully monitored clinically for signs of toxicity, and doses should be adjusted accordingly.

#### ***Hepatic impairment:***

Data on cycloserine use in hepatic impairment are scarce. Patients should be carefully monitored for signs of toxicity.

#### ***Duration of therapy***

Therapy should be continued long enough to prevent relapse.

The duration of tuberculosis therapy depends on the regimen chosen, the patient’s clinical and radiographical responses, smear and culture results, and susceptibility studies of *Mycobacterium tuberculosis* isolates from the patient or the suspected source case.

If therapy is interrupted, the treatment schedule should be extended to a later completion date depending, e.g., on the length of the interruption, the time during therapy (early or late) or the patient’s status.

#### **Method of administration**

Cycloserine is best taken without food. It can be taken with orange juice.

#### ***Extemporaneous formulation for children***

One small bowl, drinking water, a teaspoon and an oral syringe are needed for preparing the extemporaneous formulation. The following steps should be applied:

1. Open the capsule and empty its contents into the cup then add 10 mL of drinking water and dissolve by stirring gently.

2. The required portion of the mixture (see dosing table above) should be withdrawn with the syringe. *If a dose other than 5 or 10 mL is recommended, or the daily dose is to be divided, the health care provider should advise the caregiver on how to measure or divide the dose and ensure a suitable oral syringe is available for measurement.*
3. The mixture should be administered immediately to the child.
4. The withdrawn mixture may be mixed with additional liquid or additional liquid may be given after administration for masking the bitter taste.
5. Any unused mixture must be discarded.

#### **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Epilepsy.

Psychiatric disease (e.g., depression, severe anxiety, psychosis).

Concurrent use of alcohol (see section 4.5).

#### **4.4 Special warnings and precautions for use**

Before initiation of treatment, bacterial susceptibility to the drug should be established.

##### **Monitoring**

Cycloserine peak concentrations should be obtained within the first 1–2 weeks of therapy and monitored regularly during therapy. The peak concentration should be kept below 35 mcg/ml. In children, peak concentrations between 15–40 µg/mL have been recommended as appropriate.

Patients should also be given blood tests and renal and hepatic function should be monitored.

##### *Neurological and mental function*

Neuropsychiatric status should be assessed at least at monthly intervals and more frequently if neuropsychiatric symptoms develop. The most dangerous risk of cycloserine is that of suicide, so mood should be carefully watched and any undue depression or personality change observed should be immediately reported. Monitoring is particularly important when used with delamanid.

Since CNS toxicity is more common with higher doses, patients receiving more than 500 mg daily should be particularly closely observed.

[TB222 trade name] should be discontinued or the dosage reduced if the patient develops symptoms of CNS toxicity, such as convulsions, psychosis, somnolence, depression, confusion, hyperreflexia, headache, tremor, vertigo, paresis, or dysarthria. Anticonvulsant drugs or sedatives may be effective in controlling these symptoms.

##### **Hypersensitivity reactions**

The drug should be discontinued if a hypersensitivity reaction (e.g., rash, hepatitis) occurs.

##### **Vitamin B6 supplementation**

Patients should receive pyridoxine (vitamin B6) while taking cycloserine. This is especially important while breastfeeding. Adults need 100 mg or more (or 50 mg per 250 mg of cycloserine) and children should receive a dose proportionate to their weight (1–2 mg/kg/day, with a usual range of 10–50 mg/day). Supplements should also be given to breastfed infants of mothers receiving cycloserine.

##### **Renal impairment**

Cycloserine should be used very cautiously in patients with renal failure (see section 4.2).

##### *Excipients*

This medicinal product contains ponceau 4R (cochineal red A), which may cause allergic reactions. It is important to consider the contribution of excipients from all the medicines that the patient is taking.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Concurrent administration of cycloserine with ethionamide, isoniazid or alcohol potentiates the neurotoxicity of cycloserine. Additive neuropsychiatric effects may also be a concern when used with delamanid, and close monitoring is important, especially in children and adolescents.

Antacids do not affect absorption of cycloserine.

##### *Food:*

Intake with a high-fat meal has been shown to negatively affect the absorption of cycloserine (see section 5.2) and should be avoided.

#### **4.6 Fertility, pregnancy and breastfeeding**

##### *Pregnancy*

Animal data do not indicate any teratogenicity. There are no or limited amount of data from the use of cycloserine in pregnant women.

Cycloserine should be given to pregnant women only if clearly needed and when there are no suitable alternatives.

##### *Breastfeeding*

Cycloserine passes into the breast milk.

If cycloserine is required by the mother, it is not a reason to discontinue breastfeeding. Exclusively breastfed infants should be monitored if this drug is used during lactation, possibly including measurement of serum levels to rule out toxicity if there is a concern.

For vitamin B6 substitution of the infant see section 4.4.

##### *Fertility*

There are no data on the effects of [TB222 trade name] on fertility.

#### **4.7 Effects on ability to drive and use machines**

No studies on the effects of [TB222 trade name] on the ability to drive and use machines have been performed. Nevertheless, the clinical status of the patient and the adverse reaction profile of cycloserine should be borne in mind when considering the patient's ability to drive or operate machinery. Negative effects of cycloserine on the ability to drive and use machines may be synergistic with the effects of alcohol (see section 4.3).

#### **4.8 Undesirable effects**

The most frequent and most important adverse reactions of cycloserine are psychiatric and central nervous system (CNS) disorders as detailed below. CNS adverse reactions appear to be dose-related, and occur within the first 2 weeks of therapy in about 15 to 30% of patients. CNS symptoms generally disappear when the drug is discontinued.

The adverse events considered at least possibly related to the treatment are listed below by body system, organ class and absolute frequency. They are not based on adequately sized randomized controlled trials, but on published literature data generated mostly during post-approval use. Therefore, frequency data may not be available in many cases.

Frequencies are defined as very common ( $\geq 1/10$ ), common ( $\geq 1/100$ ,  $< 1/10$ ), uncommon ( $\geq 1/1000$ ,  $< 1/100$ ), rare ( $\geq 1/10,000$ ,  $< 1/1000$ ), very rare ( $\leq 1/10,000$ ), 'not known'.

### **Blood and lymphatic system disorders**

Frequency not known      Vitamin B12 deficiency, folic acid deficiency, megaloblastic anaemia, sideroblastic anaemia

### **Cardiac disorders**

Rare      Cardiac arrhythmias and sudden development of congestive heart failure in patients receiving 1 g or more per day.

### **Hepatobiliary disorders**

Not known      Elevated serum transaminases, particularly in patients with liver disease

### **Immune system disorders**

Rare      Hypersensitivity reactions including rash, photosensitivity or hepatitis

### **Nervous system disorders**

Very common      headache, tremor, dysarthria, vertigo

Not known      dysarthria, major and minor clonic seizures, convulsions, coma, paresis, hyperreflexia, paresthesia, peripheral neuropathy

### **Psychiatric disorders**

Very common      depression, confusion, anxiety, nervousness, drowsiness, dizziness, somnolence, lethargy

Not known      disorientation, loss of memory, psychoses, suicidal tendencies, aggression, character changes,

### **Skin and subcutaneous tissue disorders**

Not known      Rash, lichenoid eruptions, Stevens-Johnson syndrome

### ***Reporting of suspected adverse reactions***

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Health care providers are asked to report any suspected adverse reactions to the marketing authorisation holder, or, if available, via the national reporting system.

## **4.9 Overdose**

Acute toxicity can occur when more than 1 g is ingested by an adult. Chronic toxicity is dose related and tends to occur if more than 500 mg are administered daily. Toxicity commonly affects the central nervous system. Effects may include headache, vertigo, confusion, drowsiness, hyperirritability, paraesthesia, slurred speech and psychosis. Following ingestion of larger doses, paresis, convulsions and coma often occur.

Symptomatic and supportive therapy is recommended. Activated charcoal may be effective in reducing absorption. Cycloserine is removed by haemodialysis.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Drugs for the treatment of tuberculosis, Antibiotics

ATC code: J04AB01

#### **Properties**

Cycloserine is a broad-spectrum antibiotic that is bacteriostatic to *Mycobacterium tuberculosis* at the clinically recommended doses.

#### **Mechanism of action**

Cycloserine is an analog of the amino acid D-alanine. It interferes with peptidoglycan formation and bacterial cell wall synthesis.

## 5.2 Pharmacokinetic properties

### Pharmacokinetics of Cycloserine

Cycloserine									
<b>Absorption</b>									
Oral bioavailability	Rapid and almost completely absorbed after oral administration								
Food effect	<table border="1"> <thead> <tr> <th></th> <th>AUC<sub>(0-∞)</sub></th> <th>C<sub>max</sub></th> <th>T<sub>max</sub></th> </tr> </thead> <tbody> <tr> <td>High fat:</td> <td>No significant effect</td> <td>27% ↓</td> <td>2.75hrs</td> </tr> </tbody> </table>		AUC <sub>(0-∞)</sub>	C <sub>max</sub>	T <sub>max</sub>	High fat:	No significant effect	27% ↓	2.75hrs
		AUC <sub>(0-∞)</sub>	C <sub>max</sub>	T <sub>max</sub>					
High fat:	No significant effect	27% ↓	2.75hrs						
<b>Distribution</b>									
Plasma protein binding <i>in vitro</i>	<20%								
Tissue distribution	Widely distributed into body tissues and fluids including lungs, ascitic fluid, pleural fluid and synovial fluid, in concentrations approximately equal to plasma concentrations of the drug								
<b>Metabolism</b>									
	30-40% of dose undergoes hepatic metabolism								
Active metabolite(s)	NA*- Metabolites have not yet been identified								
<b>Elimination</b>									
Elimination half life	Between 4 and 30 hours with mean 10 hours								
% of dose excreted in urine	60 - 70% as unchanged drug, the remainder as metabolites								
% of dose excreted in faeces	Small amounts of the drug are excreted in faeces								

\*Information not available

## 5.3 Preclinical safety data

There are no additional preclinical data of relevance to the prescriber beyond those already included in other sections of the Summary of Product Characteristics.

## 6. PHARMACEUTICAL PARTICULARS

### 6.1 List of excipients

*Capsule content:*

talc

*Capsule shell composition:*

Gelatin

Ponceau 4R (cochineal red A) (E124)

quinoline yellow (E104)

titanium dioxide (E171)

### 6.2 Incompatibilities

Not applicable

### **6.3 Shelf life**

24 months

### **6.4 Special precautions for storage**

Do not store above 25°C. Store in a dry place. Keep the bottle tightly closed to protect from moisture. Protect from light.

Discard the product 50 days after initial opening.

### **6.5 Nature and contents of container**

HDPE bottle with HDPE screw cap, provided with a tamper-proof ring and induction foil sealing. 1 bottle with a leaflet in a carton. Pack size: 21 or 100 capsules.

### **6.6 Special precautions for disposal**

No special requirements.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## **7. SUPPLIER**

Biocom JSC

54 Chapaevsky Cr

355016 Stavropol

Russian Federation

Tel. No.: +78652365356

Fax No.: 82-2-924-2662

Email: [biocom@biocom.ru](mailto:biocom@biocom.ru)

## **8. WHO REFERENCE NUMBER (WHO Prequalification Programme)**

TB222

## **9. DATE OF PREQUALIFICATION**

20 August 2013

## **10. DATE OF REVISION OF THE TEXT**

March 2022

### **References**

1. WHO consolidated guidelines on tuberculosis, Module 4: Treatment. Drug-resistant tuberculosis treatment 2020  
<https://apps.who.int/iris/bitstream/handle/10665/332397/9789240007048-eng.pdf>
2. Mirzayev F, Viney K, Linh NN, et al. World Health Organization recommendations on the treatment of drug-resistant tuberculosis, 2020 update. Eur Respir J 2020; in press  
<https://doi.org/10.1183/13993003.03300-2020>
3. WHO, Geneva 2020. WHO operational handbook on tuberculosis, module 4: treatment. Drug-resistant tuberculosis treatment. <https://apps.who.int/iris/handle/10665/332398>.
4. WHO, Geneva 2010 Guidelines for treatment of tuberculosis. Fourth edition:  
<http://www.who.int/tb/publications/2010/9789241547833/en/>
5. WHO, Geneva 2011 Guidelines for the programmatic management of drug-resistant tuberculosis. update:  
<http://www.who.int/tb/publications/tb-drugresistant-guidelines/en/>
6. WHO, Geneva 2008 WHO Management of MDR-TB.A field guide,  
[http://apps.who.int/iris/bitstream/10665/44163/1/9789241547765\\_eng.pdf?ua=1&ua=1](http://apps.who.int/iris/bitstream/10665/44163/1/9789241547765_eng.pdf?ua=1&ua=1)
7. WHO, Geneva, 2014: Companion handbook to the WHO guidelines for the programmatic management of drug-resistant tuberculosis.



8. Seromycin: Package Insert and label Information, October 2011 (<https://druginserts.com/lib/rx/meds/seromycin-1/>)
9. Martindale: The Complete Drug Reference, 2017, The Pharmaceutical Press.
10. Morton RF, et al. Studies on the absorption, diffusion, and excretion of cycloserine. *Antibiot Annu* 1955-56; 3: 169-72.
11. Zhu M, et al. Pharmacokinetics of cycloserine under fasting conditions and with high-fat meal, orange juice, and antacids. *Pharmacotherapy*. 2001 (8):891-7.
12. Drugs and Lactation Database (LactMed). Available at: <https://www.ncbi.nlm.nih.gov/books/NBK501302/>
13. WHO consolidated guidelines on tuberculosis, Module 5: Management of tuberculosis in children and adolescents 2022. <https://apps.who.int/iris/rest/bitstreams/1414329/retrieve>
14. WHO, Geneva 2022. WHO operational handbook on tuberculosis. Module 5: Management of tuberculosis in children and adolescents. <https://apps.who.int/iris/rest/bitstreams/1414333/retrieve>

*Detailed information on this medicine is available on the World Health Organization (WHO) website:*  
<https://extranet.who.int/pqweb/medicines>