

WHO-PQ RECOMMENDED PATIENT INFORMATION LEAFLET

*This patient information leaflet focuses on uses of the medicine covered by WHO's Prequalification Team - Medicines. The recommendations for use are based on WHO guidelines and on information from stringent regulatory authorities.**

The medicine may be authorised for additional or different uses by national medicines regulatory authorities.

*https://extranet.who.int/pqweb/sites/default/files/documents/75%20SRA%20clarification_Feb2017_newtempl.pdf

Information for the patient

[TB226 trade name][†]

Ethambutol hydrochloride

*The warnings and instructions in this leaflet are intended for the person taking the medicine.
If you are a parent or carer responsible for giving the medicine to someone else such as a child,
you will need to apply the instructions accordingly.*

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have questions about the medicine, ask your health care provider.
- This medicine is for you only. Do not pass it on to others. It may harm them, even if their illness seems to be the same as yours.
- If you are concerned about any side effects, talk to your health care provider. This includes unwanted effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What [TB226 trade name] is and what it is used for
2. What you need to know before you take [TB226 trade name]
3. How to take [TB226 trade name]
4. Possible side effects
5. How to store [TB226 trade name]
6. Contents of the pack and other information

1. What [TB226 trade name] is and what it is used for

[TB226 trade name] is a medicine used to treat tuberculosis (TB), an infection caused by bacteria called *Mycobacterium tuberculosis*. It is always given together with other medicines to treat tuberculosis. Your healthcare provider has chosen the combination that is right for your condition.

[TB226 trade name] contains the active substance ethambutol. It works by stopping the bacteria from multiplying in the body.

2. What you need to know before you take [TB226 trade name]

Do not use [TB226 trade name] if you:

- are allergic (hypersensitive) to ethambutol, or to any of the other ingredients of this medicine (listed in section 6)

Warnings and precautions

Talk to your health care provider if you have:

- problems with eyesight. Contact your health care provider at once if you notice any changes with your vision, blurring, difficulty telling colours apart, trouble seeing, or eye pain. Your health care provider will check your vision before treatment and also during treatment.
- kidney disease. Your healthcare provider may change how you take this medicine if your kidney don't work well enough.

[†]Trade names are not prequalified by WHO. This is the national medicines regulatory agency's responsibility.

Taking other medicines

Tell your health care provider if you are taking, have recently taken or might take any other medicines.

You should not take [TB226 trade name] at the same time as stomach medicines that contain aluminium hydroxide (used to relieve heartburn and indigestion), as this can make ethambutol less effective.

Pregnancy

[TB226 trade name] can be used during pregnancy for treating tuberculosis.

Tuberculosis can be particularly dangerous in pregnancy and it is important that you get the right treatment.

Breast-feeding

You can breast-feed your baby during treatment with [TB226 trade name].

Ethambutol passes into breast milk but it is not likely to have any unwanted effects on your baby.

Driving and using machines

[TB226 trade name] may cause eyesight problems or tingling or numbness in the hands and feet. Very rarely, the medicine can also cause dizziness and disorientation. These side effects can affect how well you can drive or use machines.

After taking [TB226 trade name], you should not drive or use machines until you know how the medicine affects you.

3. How to take [TB226 trade name]

Take [TB226 trade name] exactly as your health care provider has told you. Always take [TB226 trade name] with other tuberculosis medicines. Your health care provider will choose the right combination of medicines for you.

Check with your health care provider if you are not sure.

The dose of [TB226 trade name] depends on your weight and whether you have *drug-sensitive tuberculosis* or *drug-resistant tuberculosis*. Your health care provider will decide the right dose for you and how long you should continue taking [TB226 trade name].

If you have kidney disease your health care provider may tell you to take [TB226 trade name] three times a week instead of taking it daily.

Drug susceptible tuberculosis

Children weighing less than 25 kg

Recommended doses of [TB226 trade name] for treating children with drug-susceptible tuberculosis are:

Patient's weight	Dose as 100-mg tablets
4 kg to less than 8 kg	1 tablet once daily
8 kg to less than 12 kg	2 tablets once daily
12 kg to less than 16 kg	3 tablets once daily
16 kg to less than 25 kg	4 tablets once daily

Adults and children weighing 25 kg or more

For patients weighing 25 kg or more, tablets containing higher amounts of ethambutol are more suitable.

Drug-resistant tuberculosis

Children weighing less than 24 kg

Recommended doses of [TB226 trade name] for treating children with drug-resistant tuberculosis are:

Patient's weight	Dose as 100-mg tablets
3 kg to less than 5 kg	½ tablet once daily
5 kg to less than 7 kg	1 tablet once daily
7 kg to less than 10 kg	2 tablets once daily
10 kg to less than 16 kg	3 tablets once daily
16 kg to less than 24 kg	4 tablets once daily

Adults and children weighing 24 kg or more

For patients weighing 24 kg or more, tablets containing higher amounts of ethambutol are more suitable.

If you take more [TB226 trade name] than you should

If you accidentally take more tablets than you should, tell a health care provider or get medical advice straight away. Take the medicine pack with you so the health care provider knows what you have taken.

If you forget to take [TB226 trade name] or you vomit after a dose

If you forget to take [TB226 trade name] at the usual time and:

- it is less than 6 hours since the usual time you take the dose, take the dose straight away and then take the next one at the usual time the next day
- if it is more than 6 hours since the usual time you take the dose, skip the missed dose and take the next one at the usual time the next day.

Do not take a double dose to make up for the forgotten dose. If you are unsure about what to do, ask your health care provider.

If you vomit within 1 hour of taking [TB226 trade name], then take another dose. If you vomit more than an hour after taking the dose, then you do not need to take another dose and you can take the next dose as usual on the next day.

If you stop taking [TB226 trade name]

It is important that you complete the course of treatment even if you begin to feel better. If you stop taking [TB226 trade name] too soon, your infection may not be completely cured and the infection may return or your condition may get worse. The bacteria causing the infection may become resistant to [TB226 trade name].

Taking this medicine

You can take [TB226 trade name] with food or between meals. Taking it with food may reduce effects on the stomach. Try to take the tablet(s) at approximately the same time each day.

[TB226 trade name] should be swallowed with water.

[TB226 trade name] may be mixed with water if the patient cannot swallow tablets.

1. Mix each tablet with at least 10 mL water but do not use more than 50 mL of water altogether.
2. Stir until the tablets break up completely and are fully mixed with the water.
3. Drink the entire medicine mixture immediately.
4. Rinse the cup with a little more water and drink this water.

If you have any questions on the use of this medicine, ask your health care provider.

4. Possible side effects

Like all medicines, [TB226 trade name] can cause side effects, but not everybody gets them. It is important that you tell the health care provider of any change in your health.

You should ask for medical help immediately if you have:

- sudden problems with breathing, wheeziness, swelling of the face, eyes, lips or throat, flushed appearance or irregular heartbeat. These may be symptoms of a severe allergic reactions

Speak with your health care provider straightaway if you have:

- changes in vision, including blurred vision, eye pain or not being able to see colours properly (colour blindness)

The following side effects can also occur in patients treated with ethambutol:

Uncommon side effects (may affect more than 1 in 1000 people)

high levels of uric acid in the blood (hyperuricaemia) which may lead to gout (pain or swelling of the joints)

Rare side effects (may affect up to 1 in 10 000 people)

tingling, 'pins-and-needles' and numbness

rash, itchiness, hives (red, itchy bumps on the skin)

blood tests showing reduced blood platelets (which can cause bleeding and bruising)

Very rare side effects (may affect fewer than 1 in 10 000 people)

dizziness, headache and disorientation

mental confusion, hallucination (seeing or hearing things that are not there)

liver problems (hepatic failure)

kidney problems (lower back pain, pain or burning when passing urine, difficulty passing urine)

feeling ill, joint pains, fever

lung damage causing breathing difficulty

skin allergy to bright light, severe skin rashes with blistering and skin peeling

blood tests showing reduced levels of some types of white blood cells

Frequency of side effects not known

nausea (feeling sick), vomiting, loss of appetite, gas (flatulence), belly pain, diarrhoea, upset stomach

inflammation of the liver, jaundice (yellowing of the skin and/or eyes), blood tests suggesting liver problems

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, tell your healthcare provider as soon as possible.

Reporting of side effects

If you get a side effect, talk to your health care provider. This includes side effects not listed in this leaflet. You may also be able to report such effects directly to your national reporting system if one is available. By reporting side effects, you can help to improve the available information on this medicine.

5. How to store [TB226 trade name]

Do not store above 30°C. Protect from light. Store in the original container. Do not remove the desiccant canister from the bottle.

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date stated on the carton or bottle after EXP. The expiry date refers to the last day of that month.

Do not use this medicine if you notice description of the visible signs of deterioration that it is different from the description below.

Do not throw away any medicines in wastewater or household waste. Ask your health care provider how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What [TB226 trade name] contains

- The active ingredient is ethambutol hydrochloride
- The other ingredients of [TB226 trade name] are:
Tablet core: microcrystalline cellulose, povidone, stearic acid, maize starch, sodium starch glycolate, colloidal anhydrous silica, purified talc and magnesium stearate
Film coat: hypromellose, ethylcellulose, macrogol, purified talc and titanium dioxide

There is too little sodium in this medicine to have any effect, even if you are on a low-sodium diet.

What [TB226 trade name] looks like and contents of the pack

[TB226 trade name] is a white, circular, shallow, biconvex, film-coated tablet, with a break-line on one side and plain on the other side.

[TB226 trade name] can be divided into equal doses.

[TB226 trade name] is available in:

- Self-sealing polythene bag, inside a 650-mL HDPE container, sealed with an aluminium tagger. Pack size: 1000 tablets.
- Dark amber coloured PVC/PVdC-Alu blisters. Pack sizes: 7 x 10, 10 x 10 and 24 x 28 tablets.

Supplier and Manufacturer

Supplier

Macleods Pharmaceuticals Limited
Atlanta Arcade, 3rd Floor, Church Road
Near Leela Hotel,
Andheri-Kurla Road
Andheri (East)
400 059 Mumbai
India
Tel. No.: +91 22-61132900
Fax No.: +91 22-28304641

Email: vijay@macleodspharma.com
sjadhav@macleodspharma.com
exports@macleodspharma.com
vijay@macleodspharma.com

Manufacturer

Macleods Pharmaceuticals Limited
Phase II/ Phase III, Unit II, Plot No. 25-27
Sr. No. 366, Premier Industrial Estate
Kachigam, 396 210 Daman
India
Tel. No.: + 91 0260 2244337 and + 91 0260 2240125
Fax No.: + 91 0260 2241565
Email: nishata@macleodspharma.com.

Oxalis Labs
Village Theda, P.O. Lodhimajra
Tehsil: Baddi
Dist. Solan
Himachal Pradesh
174101, India
Tel. No.: +91-1795661400
Fax No.: +91-1795661452

For any information about this medicine, contact the supplier.

This leaflet was last revised in November 2023

Detailed information on this medicine is available on the World Health Organization (WHO) website: <https://extranet.who.int/pqweb/medicines>