

# Slovakia

Population 2016

5.4 million

Estimates of TB burden*, 2016	Number (thousands)	Rate (per 100 000 population)
Mortality (excludes HIV+TB)	0.035 (0.034–0.035)	0.64 (0.63–0.65)
Mortality (HIV+TB only)	<0.01 (<0.01–<0.01)	0 (0–0)
Incidence (includes HIV+TB)	0.32 (0.28–0.37)	5.9 (5.1–6.9)
Incidence (HIV+TB only)	0 (0–0)	0.01 (0–0.01)
Incidence (MDR/RR-TB)**	<0.01 (0–<0.01)	0.03 (0.01–0.05)

Estimated TB incidence by age and sex (thousands)*, 2016			
	0-14 years	> 14 years	Total
Females	0.015 (0.013–0.018)	0.095 (0.081–0.11)	0.11 (0.094–0.13)
Males	0.016 (0.014–0.018)	0.2 (0.17–0.23)	0.21 (0.18–0.24)
Total	0.031 (0.027–0.036)	0.29 (0.25–0.34)	0.32 (0.28–0.37)

TB case notifications, 2016	
Total cases notified	296
Total new and relapse	281
- % tested with rapid diagnostics at time of diagnosis	21%
- % with known HIV status	0%
- % pulmonary	85%
- % bacteriologically confirmed among pulmonary	60%

Universal health coverage and social protection	
TB treatment coverage (notified/estimated incidence), 2016	87% (75–100)
TB patients facing catastrophic total costs	
TB case fatality ratio (estimated mortality/estimated incidence), 2016	0.11 (0.09–0.13)

TB/HIV care in new and relapse TB patients, 2016		Number	(%)
Patients with known HIV-status who are HIV-positive		0	
- on antiretroviral therapy		0	

Drug-resistant TB care, 2016			
	New cases	Previously treated cases	Total number***
Estimated MDR/RR-TB cases among notified pulmonary TB cases			1 (0–4)
Estimated % of TB cases with MDR/RR-TB	0% (0–2.6)	3.7% (0.09–19)	
% notified tested for rifampicin resistance	37%	41%	111
MDR/RR-TB cases tested for resistance to second-line drugs			7
Laboratory-confirmed cases		MDR/RR-TB: 7, XDR-TB: 1	
Patients started on treatment ****		MDR/RR-TB: 2, XDR-TB: 1	

Treatment success rate and cohort size		
	Success	Cohort
New and relapse cases registered in 2015	86%	305
Previously treated cases, excluding relapse, registered in 2015	78%	9
HIV-positive TB cases registered in 2015		0
MDR/RR-TB cases started on second-line treatment in 2014	50%	4
XDR-TB cases started on second-line treatment in 2014		0

TB preventive treatment, 2016	
% of HIV-positive people (newly enrolled in care) on preventive treatment	
% of children (aged < 5) household contacts of bacteriologically-confirmed TB cases on preventive treatment	100% (100–100)

TB financing, 2017	
National TB budget (US\$ millions)	<1


\* Ranges represent uncertainty intervals

\*\* MDR is TB resistant to rifampicin and isoniazid; RR is TB resistant to rifampicin


\*\*\* Includes cases with unknown previous TB treatment history

\*\*\*\* Includes patients diagnosed before 2016 and patients who were not laboratory-confirmed

## Tuberculosis profile





— Mortality (excludes HIV+TB)



— Incidence  
— Notified (new and relapse)  
— Incidence (HIV+TB only)

Notified cases by age group and sex, 2016



— New and relapse  
— Retreatment, excluding relapse  
— HIV-positive — MDR/RR-TB — XDR-TB